

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problems Mailbox.**

This Page Blank (uspto)

12

EUROPEAN PATENT APPLICATION

21 Application number: **87400544.0**

51 Int. Cl. 4: **C 12 N 15/00**

22 Date of filing: **11.03.87**

A 01 H 1/00, C 12 N 5/00,
A 01 H 5/10
// C12N9/10

30 Priority: **11.03.86 EP 86400521**
21.01.87 EP 87400141

43 Date of publication of application:
21.10.87 Bulletin 87/43

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

71 Applicant: **PLANT GENETIC SYSTEMS N.V.**
Kunstlaan Avenue des Arts, 46
B-1040 Bruxelles (BE)

BIOGEN N.V.
15 Pietermaai
Willemstad Curaçao Netherlands Antilles (NL)

72 Inventor: **Leemans, Jan**
Ellebochte 38
B-9210 Heusden (BE)

Botterman, Johan
Eedstraat 37
B-9710 Zwijnaarde (BE)

de Block, Marc
Vogelhoeckstraat 32
B-9219 Gentbrugge (BE)

Thompson, Charles
19 chemin des Palettes
CH-1212 Grand Lancy/Geneve (CH)

Mouva, Rao
10, avenue Secheron
CH-1202 Geneve (CH)

74 Representative: **Gutmann, Ernest et al**
S.C. Ernest Gutmann - Yves Plasseraud 67, boulevard
Haussmann
F-75008 Paris (FR)

The title of the invention has been amended (Guidelines for Examination in the EPO, A-III, 7.3).
The microorganism(s) has (have) been deposited at The German Collection of Micro-Organisms (DSM) under number(s) DSM 3606, DSM 3687, DSM 4023.

54 **Plant cells resistant to glutamine synthetase inhibitors, made by genetic engineering.**

57 The invention relates to a DNA fragment containing a determined gene, the expression of which inhibits the antibiotic and herbicidal effects of Bialaphos and related products.

It also relates to recombinant vectors, containing such DNA fragment, which enable this protective gene to be introduced and expressed into cells and plant cells.

Description

GENETICALLY ENGINEERED PLANT CELLS AND PLANTS EXHIBITING RESISTANCE TO GLUTAMINE SYNTHETASE INHIBITORS, DNA FRAGMENTS AND RECOMBINANTS FOR USE IN THE PRODUCTION OF SAID CELLS AND PLANTS.

The invention relates to a process for protecting plant cells and plants against the action of glutamine synthetase inhibitors.

It also relates to applications of such process, particularly to the development of herbicide resistance into determined plants.

It relates further to non-biologically transformed plant cells and plants displaying resistance to glutamine synthetase inhibitors as well as to suitable DNA fragments and recombinants containing nucleotide sequences encoding resistance to glutamine synthetase inhibitors.


Glutamine synthetase (hereafter simply designated by GS) constitutes in most plants one of the essential enzymes for the development and life of plant cells. It is known that GS converts glutamate into glutamine. GS is involved in an efficient pathway (the only one known nowadays) in most plants for the detoxification of ammonia released by nitrate reduction, amino acid degradation or photorespiration. Therefore potent inhibitors of GS are very toxic to plant cells. A particular class of herbicides has been developed, based on the toxic effect due to inhibit inhibition of GS in plants.

These herbicides comprise as active ingredient a GS inhibitor.


There are at least two possible ways which might lead to plants resistant to the inhibitors of the action of glutamine synthetase; (1) by changing the target. It can be envisaged that mutations in the GS enzyme can lead to insensitivity towards the herbicide; (2) by inactivation of the herbicide. Breakdown or modification of the herbicide inside the plant could lead to resistance.

Bialaphos and phosphinothricin (hereafter simply designated by PPT) are two such inhibitors of the action of GS, (ref. 16, 17) and have been shown to possess excellent herbicidal properties (see more particularly ref. 2 as concerns Bialaphos).

Bialaphos has the following formula (I) :



PPT has the following formula (II) :



Thus the structural difference between PPT and Bialaphos resides in the absence of two alanine amino acids in the case of PPT.

These two herbicides are non selective. They inhibit growth of all the different species of plants present on the soil, accordingly cause their total destruction.

Bialaphos was first disclosed as having antibiotic properties, which enabled it to be used as a pesticide or a fungicide. Bialaphos can be produced according to the process disclosed in the United States patent no 3 832 394, assigned to MEIJI SEIKA KAISHA LTD., which patent is incorporated herein by reference. It comprises cultivating *Streptomyces hygroscopicus*, such as the strain available at the American Type Culture Collection, under the ATCC number 21,705, and recovering Bialaphos from its culture medium. However, other strains, such as *Streptomyces viridochromogenes*, also produce this compound (ref. 1).

Other tripeptide antibiotics which contain a PPT moiety are or might be discovered in nature as well, e.g. phosalacin (ref. 15).

PPT is also obtained by chemical synthesis and is commercially distributed by the industrial Company HOECHST.

A number of *Streptomyces* species have been disclosed which produce highly active antibiotics which are known to incapacitate prokaryotic cell functions or enzymes. The *Streptomyces* species which produce these antibiotics would themselves be destroyed if they had not a self defence mechanism against these antibiotics. This self defence mechanism has been found in several instances to comprise an enzyme capable of inhibiting the antibiotic effect, thus of avoiding autotoxicity for the *Streptomyces* species concerned. This modification is generally reversed when the molecule is exported from the cell.

The existence of a gene which encodes an enzyme able to modify the antibiotic so as to inhibit the antibiotic effect against the host has been demonstrated in several *Streptomyces* producing antibiotics, for example, in *S. fradiae*, *S. azureus*, *S. vinaceus*, *S. erythreus*, producing neomycin, thiostrepton, viomycin, and MLS (Macrolide Lincosamide Streptogramin) antibiotics respectively (ref. 4), (ref. 5), (ref. 6), (ref. 14 by CHATER et al., 1982 describes standard techniques which can be used for bringing these effects to light).

In accordance with the present invention, it has been found that *Streptomyces hygroscopicus* ATCC 21,705, also possesses a gene encoding an enzyme responsible of the inactivation of the antibiotic properties of Bialaphos. Experiments carried out by the applicants have lead to the isolation of such a gene and its use in a process for controlling the action of GS inhibitors, based on PPT or derived products.

An object of the invention is to provide a new process for controlling the action in plant cells and plants of GS inhibitors.

Another object of the invention is to provide DNA fragments and DNA recombinants, particularly modified vectors containing said DNA fragments, which DNA fragments contain nucleotide sequences capable, when incorporated in plant cells and plants, to protect them against the action of GS inhibitors.

A further object of the invention is to provide non-biologically transformed plant cells and plants capable of neutralizing or inactivating GS inhibitors.

A further object of the invention is to provide a process for selectively protecting plant species against herbicides of a GS inhibitor type.

More specifically an object of the invention is to provide a DNA fragment transferable to plant cells- and to whole plants- capable of protecting them against the herbicidal effects of Bialaphos and of structurally analogous herbicides.

A further object of the invention is to provide plant cells resistant to the products of the class exemplified by Bialaphos, which products possess the PPT unit in their structure.

The process according to the invention for controlling the action in plant cells and plants of a GS inhibitor when contacted therewith, comprises providing said plants with a heterologous DNA fragment including a foreign nucleotide sequence, capable of being expressed in the form of a protein in said plant cells and plants, under condition such as to cause said heterologous DNA fragment to be integrated stably through generations in the cells of said plants, and wherein said protein has an enzymatic activity capable of inactivating or neutralization of said glutamine synthetase inhibitor.

A preferred DNA fragment is one derived from an antibiotic-producing-*Streptomyces* strain (or a sequence comprising a nucleotide sequence encoding the same activity) and which encodes resistance to a said GS inhibitors.

Preferred nucleotide sequences for use in this invention encode a protein which has acetyl tranferase activity with respect to said GS inhibitors.

A most preferred DNA fragment according to the invention comprises a nucleotide sequence coding for a polypeptide having a PPT acetyl transferase activity.

A particular DNA fragment according to the invention, for the subsequent transformation of plant cells, consists of a nucleotide sequence coding for at least part of a polypeptide having the following sequence :

X SER PRO GLU

183
 5 ARG ARG PRO ALA ASP ILE ARG ARG ALA THR GLU ALA ASP MET PRO
 228
 ALA VAL CYS THR ILE VAL ASN HIS TYR ILE GLU THR SER THR VAL
 10 273
 ASN PHE ARG THR GLU PRO GLN GLU PRO GLN GLU TRP THR ASP ASP
 318
 15 LEU VAL ARG LEU ARG GLU ARG TYR PRO TRP LEU VAL ALA GLU VAL
 363
 ASP GLY GLU VAL ALA GLY ILE ALA TYR ALA GLY PRO TRP LYS ALA
 20 408
 ARG ASN ALA TYR ASP TRP THR ALA GLU SER THR VAL TYR VAL SER
 453
 25 PRO ARG HIS GLN ARG THR GLY LEU GLY SER THR LEU TYR THR HIS
 498
 LEU LEU LYS SER LEU GLU ALA GLN GLY PHE LYS SER VAL VAL ALA
 30 543
 VAL ILE GLY LEU PRO ASN ASP PRO SER VAL ARG MET HIS GLU ALA
 588
 35 LEU GLY TYR ALA PRO ARG GLY MET LEU ARG ALA ALA GLY PHE LYS
 633
 HIS GLY ASN TRP HIS ASP VAL GLY PHE TRP GLN LEU ASP PHE SER
 40 678
 LEU PRO VAL PRO PRO ARG PRO VAL LEU PRO VAL THR GLU ILE
 723

45
 in which X represents MET or VAL, which part of said polypeptide is of sufficient length to confer protection against Bialaphos to plant cells, when incorporated genetically and expressed therein, i.e. as termed hereafter "plant-protecting capability" against Bialaphos.

50 A preferred DNA fragment consists of the following nucleotide sequence :

55

60

65

183 GTG AGC CCA GAA
 CGA CGC CCG GCC GAC ATC CGC CGT GCC ACC GAG GCG GAC ATG CCG
 228
 GCG GTC TGC ACC ATC GTC AAC CAC TAC ATC GAG ACA AGC ACG GTC
 273
 AAC TTC CGT ACC GAG CCG CAG GAA CCG CAG GAG TGG ACG GAC GAC
 318
 CTC GTC CGT CTG CCG GAG CGC TAT CCC TGG CTC GTC GCC GAG GTG
 363
 GAC GGC GAG GTC GCC GGC ATC GCC TAC GCG GGC CCC TGG AAG GCA
 408
 CCG AAC GCC TAC GAC TGG ACG GCC GAG TCG ACC GTG TAC GTC TCC
 453
 CCC CGC CAC CAG CCG ACG GGA CTG GGC TCC ACG CTC TAC ACC CAC
 498
 CTG CTG AAG TCC CTG GAG GCA CAG GGC TTC AAG AGC GTG GTC GCT
 543
 GTC ATC GGG CTG CCC AAC GAC CCG AGC GTG CGC ATG CAC GAG GCG
 588
 CTC GGA TAT GCC CCC CGC GGC ATG CTG CCG GCG GCC GGC TTC AAG
 633
 CAC GGG AAC TGG CAT GAC GTG GGT TTC TGG CAG CTG GAC TTC AGC
 678
 CTG CCG GTA CCG CCC CGT CCG GTC CTG CCC GTC ACC GAG ATC
 723



or of a part thereof expressing a polypeptide having plant protecting capability against Bialaphos.
 The invention also relates to any DNA fragment differing from the preferred one indicated hereabove by the replacement of any of its nucleotides by others, yet without modifying the genetic information of the preferred DNA sequence mentioned hereabove (normally within the meaning of the universal genetic code), and furthermore to any equivalent DNA sequence which would encode a polypeptide having the same properties, particularly a Bialaphos-resistance-activity.

It will be understood that the man skilled in the art should be capable of readily assessing those parts of the nucleotide sequences that could be removed from either side of any of the DNA fragments according to the invention, for instance by removing terminal parts on either side of said DNA fragment, such as by an exonucleolytic enzyme, for instance Bal31, by recloning the remaining fragment in a suitable plasmid and by assaying the capacity of the modified plasmid to transform appropriate cells and to protect it against the Bialaphos antibiotic or herbicide as disclosed later, whichever assay is appropriate.

For the easiness of language, these DNA fragments will be termed hereafter as "Bialaphos-resistance DNA". In a similar manner, the corresponding polypeptide will be termed as "Bialaphos-resistance enzyme".

While in the preceding discussion particular emphasis has been put on DNA fragments capable, when introduced into plant cells and plants, to confer on them protection against Bialaphos or PPT, it should be understood that the invention should in no way be deemed as limited thereto.

In a same manner, the invention pertains to DNA fragments which, when introduced into such plant cells, would also confer on them a protection against other GS inhibitors, for instance of intermediate products involved in the natural biosynthesis of phosphinotricin, such as the compounds designated by the abbreviations MP101 (III), MP102 (IV), the formula of which are indicated hereafter :



More generally, the invention has opened the route to the production of DNA fragments which, upon proper incorporation into plant cells and plants, can protect them against GS inhibitors when contacted therewith, as this will be shown in a detailed manner in relation to Bialaphos and PPT in the examples which will follow.

This having been established, it will be appreciated that any fragment encoding an enzymatic activity which would protect plant cells and plants against said GS inhibitors, by inactivation, should be viewed as an equivalent of the preferred fragments which have been disclosed hereabove. This would apply especially to any DNA fragments that would result from genetic screening of the genomic DNAs of strains, particularly of antibiotic-producing strains, likely to possess genes which, even though structurally different, would encode similar activity with respect to Bialaphos or PPT, or even with respect to other GS inhibitors. This applies to any gene in other strains producing a PPT derivative.

Therefore, it should be understood that the language "Bialaphos-resistance DNA" or "Bialaphos-resistance enzyme" used thereafter as a matter of convenience is intended to relate not only to the DNAs and enzymes specifically concerned with resistance to PPT or most directly related derivatives, but more generally with other DNAs and enzymes which would be capable, under the same circumstances, of inactivating the action in plants of GS inhibitors.

The invention also relates to DNA recombinants containing the above defined Bialaphos-resistance DNA fragments recombined with heterologous DNA, said heterologous DNA containing regulation elements and said Bialaphos-resistance DNA being under the control of said regulation elements in such manner as to be expressible in a foreign cellular environment compatible with said regulation elements. Particularly the abovesaid Bialaphos-resistance-DNA fragments contained in said DNA recombinants are devoid of any DNA region involved in the biosynthesis of Bialaphos, when said Bialaphos-resistance-DNA fragment originate themselves from Bialaphos-producing strains.

By "heterologous DNA" is meant a DNA of an other origin than that from which said Bialaphos-resistance-DNA originated, e.g. is different from that of a *Streptomyces hygroscopicus* or *Streptomyces viridochromogenes* or even more preferably a DNA foreign to *Streptomyces* DNA. Particularly said regulation elements are those which are capable of controlling the transcription and translation of DNA sequences normally associated with them in said foreign environment. "Cellular" refers both to micro-organisms and to cell cultures.

This heterologous DNA may be a bacterial DNA, particularly when it is desired to produce a large amount of the recombinant DNA, such as for amplification purposes. In that respect a preferred heterologous DNA consists of DNA of *E. coli* or of DNA compatible with *E. coli*. It may be DNA of the same origin as that of the cells concerned or other DNA, for instance viral or plasmidic DNA known as capable of replicating in the cells concerned.

Preferred recombinant DNA contains heterologous DNA compatible with plant cells, particularly Ti-plasmid DNA.

Particularly preferred recombinants are those which contain GS inhibitor inactivating DNA under the control of a promoter recognized by plant cells, particularly those plant cells on which inactivation of GS inhibitors is to be conferred.

Preferred recombinants according to the invention further relate to modified vectors, particularly plasmids, containing said GS-inhibitor-inactivating DNA so positioned with respect to regulation elements, including particularly promoter elements, that they enable said GS inhibitor-inactivating DNA to be transcribed and translated in the cellular environment which is compatible with said heterologous DNA. Advantageous vectors are those so engineered as to cause stable incorporation of said GS inhibitor inactivating DNA in foreign cells, particularly in their genomic DNA. Preferred modified vectors are those which enable the stable transformation

of plant cells and which confer to the corresponding cells, the capability of inactivating GS inhibitors. It seems that, as described later, the initiation codon of the Bialaphos-resistance-gene of the *Streptomyces hygroscopicus* strain used herein is a GTG codon. But in preferred recombinant DNAs or vectors, the Bialaphos-resistance-gene is modified by substitution of an ATG initiation codon for the initiation codon GTG, which ATG enables translation initiation in plant cells.

In the example which follows, the plant promoter sequence which has been used was constituted by a promoter of the 35 S cauliflower mosaic virus. Needless to say that the man skilled in the art will be capable of selecting other plant promoters, when more appropriate in relation to the plant species concerned.

According to an other preferred embodiment of the invention, particularly when it is desired to achieve transport of the enzyme encoded by the Bialaphos-resistance-DNA into the chloroplasts, the heterologous DNA fragment is fused to a gene or a DNA fragment encoding a transit peptide, said last mentioned fragment being then intercalated between the GS inhibitor inactivating gene and the plant promoter selected.

As concerns means capable of achieving such constructions, reference can be made to the following British applications 84 32757 filed on December 28, 1984 and 85 00336 filed on January 7, 1985 and to the related applications filed in the United-States of America (no 755,173, filed on July 15, 1985), in the European Patent Office (no 85 402596.2, filed on December 20, 1985), in Japan (no 299 730, filed on December 27, 1985), in Israel (no 77 466 filed on December 27, 1985) and in Australia (no 5 165 485, filed on December 24, 1985), all of which are incorporated herein by reference.

Reference can also be made to the scientific literature, particularly to the following articles :
- VAN DEN BROECK et al., 1985, Nature, 313, 358-363 ;
- SCHREIER and al., Embo. J., vol. 4, no 1, 25-32.

These articles are also incorporated herein by reference.

For the sake of the record, be it recalled here that under the expression "transit peptide", one refers to a polypeptide fragment which is normally associated with a chloroplast protein or a chloroplast protein sub-unit in a precursor protein encoded by plant cell nuclear DNA. The transit peptide then separates from the chloroplast protein or is proteolitically removed, during the translocation process of the latter protein into the chloroplasts. Examples of suitable transit peptides are those associated with the small subunit of ribulose-1,5 biphosphate (RuBP) carboxylase or that associated with the chlorophyll a/b binding proteins.

There is thus provided DNA fragments and DNA recombinants which are suitable for use in the process defined hereafter.

More particularly the invention also relates to a process, which can be generally defined as a process for producing plants and reproduction material of said plants including a heterologous genetic material stably integrated therein and capable of being expressed in said plants or reproduction material in the form of a protein capable of inactivating or neutralizing the activity of a glutamine synthetase-inhibitor, comprising the non biological steps of producing plants cells or plant tissue including said heterologous genetic material from starting plant cells or plant tissue not able to express that inhibiting or neutralizing activity, regenerating plants or reproduction material of said plants or both from said plant cells or plant tissue including said genetic material and, optionally, biologically replicating said last mentioned plants or reproduction material or both, wherein said non-biological steps of producing said plant cells or plant tissue including said heterologous genetic material, comprises transforming said starting plant cells or plant tissue with a DNA-recombinant containing a nucleotide sequence encoding said protein, as well as the regulatory elements selected among those which are capable of enabling the expression of said nucleotide sequence in said plant cells or plant tissue, and to cause the stable integration of said nucleotide sequence in said plant cells and tissue, as well as in the plant and reproduction material processed therefrom throughout generations.

The invention also relates to the cell cultures containing Bialaphos-resistance-DNA, or more generally said GS-inhibitor-inactivating DNA, which cell cultures have the property of being resistant to a composition containing a GS inhibitor, when cultured in a medium containing a such composition at dosages which would be destructive for non transformed cells.

The invention concerns more particularly those plant cells or cell cultures in which the Bialaphos-resistance DNA is stably integrated and which remains present over successive generations of said plant cells. Thus the resistance to a GS inhibitor, more particularly Bialaphos or PPT, can also be considered as a way of characterizing the plant cells of this invention.

Optionally one may also resort to hybridization experiments between the genomic DNA obtained from said plant cells with a probe containing a GS inhibitor inactivating DNA sequence.

More generally the invention relates to plant cells, reproduction material, particularly seeds, as well as plants containing a foreign or heterologous DNA fragment stably integrated in their respective genomic DNAs, said fragments being transferred throughout generations of such plant cells, reproduction material, seeds and plants, wherein said DNA fragment encodes a protein inducing a non-variety-specific enzymatic activity capable of inactivating or neutralizing GS inhibitors, particularly Bialaphos and PPT, more particularly to confer on said plant cells, reproduction material, seeds and plants a corresponding non-variety-specific phenotype of resistance to GS inhibitors.

"Non-variety-specific" enzymatic activity or phenotype aims at referring to the fact that they are not characteristic of specific plant genes or species as this will be illustrated in a non-limitative way by the examples which will follow. They are induced in said plant materials by essentially non-biological processes applicable to plants belonging to species normally unrelated with one another and comprising the

incorporation into said plant material of heterologous DNA, e.g. bacterial DNA or chemically synthesized DNA, which does not normally occur in said plant material or which normally cannot be incorporated therein by natural breeding processes, and which yet confers a common phenotype (e.g. herbicide resistance) to them.

The invention is of particular advantageous use in processes for protecting field-cultivated plant species against weeds, which processes comprise the step of treating the field with an herbicide, e.g. Bialaphos or PPT in a dosage effective to kill said weeds, wherein the cultivated plant species then contains in their genome a DNA fragment encoding a protein having an enzymatic activity capable of neutralizing or inactivating said GS inhibitor.

By way of illustration only, effective doses for use in the abovesaid process range from about 0.4 to about 1.6 kg/Hectare of Bialaphos or PPT.

There follows now a disclosure of how the preferred DNA fragment described hereabove was isolated starting from the *Streptomyces hygroscopicus* strain available at the American Type Culture Collection under deposition number ATCC 21 705, by way of exemplification only.

The following disclosure also provides the technique which can be applied to other strains producing compounds with a PPT moiety.

The disclosure will then be completed with the description of the insertion of a preferred DNA fragment conferring to the transformed cells the capability of inactivating Bialaphos and PPT. Thus the Bialaphos-inactivating-DNA fragment designated thereafter by Bialaphos-resistance gene or "sfr" gene, isolated by the above described technique into plasmids which can be used for transforming plant cells and conferring to them a resistance against Bialaphos, also merely by way of example for non-limitative illustration purposes.

The following disclosure is made with reference to the drawings in which :

- fig. 1 is a restriction map of a plasmid containing a *Streptomyces hygroscopicus* DNA fragment encoding Bialaphos-resistance, which plasmid, designated hereafter as pBG1 has been constructed according to the disclosure which follows ;

- fig. 2 shows the nucleotide sequence of smaller fragment obtained from pBG1, subcloned into another plasmid (pBG39) and containing the resistance gene ;

- fig. 3 shows the construction of a series of plasmids given by way of example, which plasmids aim at providing suitable adaptation means for the insertion therein of the Bialaphos-resistance gene or "sfr" gene ;

- fig. 4A and 4B show the construction of a series of plasmids given by way of example, which plasmids contain suitable plant cell promoter sequences able to initiate transcription and expression of the foreign gene inserted under their control into said plasmids ;

- fig. 5A shows a determined fragment of the nucleotide sequence of the plasmid obtained in figure 3 ;

- fig. 5B shows the reconstruction of the first codons of a Bialaphos-resistance gene, from a *FokI*/*Bgl*III fragment obtained from pBG39 and the substitution of an ATG initiation codon for the GTG initiation codon of the natural "sfr" gene ;

- fig. 5C shows the reconstruction of the entire "sfr" gene, namely the last codons thereof, and its insertion into a plasmid obtained in figures 4A and 4B ;

- fig. 6A shows an expression vector containing the "sfr" gene placed under the control of a plant cell promoter ;

- fig. 6B shows another expression vector deriving from the one shown in fig. 6A, by the substitution of some nucleotides.

- fig. 7 shows the construction of a series of plasmids given by way of examples, to ultimately produce plasmids containing the promoter region and the transit peptide sequence of a determined plant cell gene, for the insertion of the "sfr" gene under the control of said promoter region and the downstream of said transit peptide sequence.

- fig. 8 to 11 will be referred to hereafter.

The following experiment was set up to isolate a Bialaphos-resistance-gene from *S. hygroscopicus*, according to standard techniques for cloning into *Streptomyces*.

2.5 µg of *S. hygroscopicus* genomic DNA and 0.5 µg of *Streptomyces* vector pIJ61 were cleaved with *Pst*I according to the method described in ref. 6. The vector fragments and genomic fragments were mixed and ligated (4 hours at 10°C followed by 72 hours at 4°C in ligation salts which contain 66 mM Tris-HCl (pH 7.5), 1 mM EDTA, 10 mM MgCl₂, 10 mM 2-mercaptoethanol and 0.1 mM ATP) at a total DNA concentration of 40 µg ml⁻¹ with T4 DNA ligase. Ligation products were introduced into 3 x 10⁹ *S. lividans* strain 66 protoplasts by a transformation procedure mediated by polyethylene-glycol (PEG) as described hereafter. These protoplasts gave rise to 5 x 10⁷ colonies and 4 x 10⁴ pocks after regeneration on 20 plates of R2 agar containing 0.5 % of Difco yeast extract (R2 YE). Preparation and composition of the different mediums and buffers used in the disclosed experiments are described hereinafter. When these lawns were replica-plated on minimal medium plates containing 50 µg ml⁻¹ Bialaphos, drug resistant colonies appeared at a frequency of 1 per 10⁴ transformants. After purification of the drug resistant colonies, there plasmid DNA was isolated and used to retransform *S. lividans* protoplasts. Non selective regeneration followed by replication to Bialaphos-containing-medium demonstrated a 100 % correlation between pocks and Bialaphos resistant growth. The recombinant plasmids of several resistant clones all contained a 1.7 Kb *Pst*I insert (see fig. 1).

Subcloning of the herbicide resistance gene

The 1.7 Kb *Pst*I insert was then subcloned into the high copy number streptomycete vector pIJ385 to generate plasmid pBG20. *S. lividans* strains which contained pBG20 were more than 500 times more resistant to Bialaphos. *S. lividans* growth is normally inhibited in minimal medium containing 1 µg/ml Bialaphos; growth of transformants containing pBG20 was not noticeably inhibited in a medium containing 500 µg/ml Bialaphos. The *Pst*I fragment was also subcloned in either orientation into the *Pst*I site of the plasmid pBR322, to produce plasmids pBG1 and pBG2, according to their orientation. A test on minimal M9 medium demonstrated that *E. coli* E8767 containing pBG1 or pBG2 was resistant to Bialaphos.

A \pm 1.65 Kb *Pst*I - *Bam*HI fragment was subcloned from pBG1 into the plasmid pUC19 to produce the plasmid pBG39, and conferred Bialaphos resistance to *E. coli*, W3110, C600 and JM83.

Using an *in vitro* coupled transcription-translation system (ref. 5) from *S. lividans* extracts, the 1.65 Kb *Pst*I - *Bam*HI fragment in pBG39 was shown to direct the synthesis of a 22 Kd protein. In the following, this 1.65 Kb insert includes a fragment coding for a 22 Kd protein and will be called "sfr" gene.

Fine mapping and sequencing of the gene

A 625 bp *Sau*3A fragment was subcloned from pBG39 into pUC19 and still conferred Bialaphos resistance to a *E. coli* W3110 host. The resulting clones were pBG93 and pBG94, according to the orientation.

The orientation of the gene in the *Sau*3A fragment was indicated by experiments which have shown that Bialaphos resistance could be induced with IPTG from the pUC19 lac promoter in pBG93. In the presence of IPTG (0.5 mM) the resistance of pBG93/W3110 increased from 5 to 50 µg/ml on a M9 medium containing Bialaphos. The W3110 host devoid of pBG93, did not grow on M9 medium containing 5 µg/ml Bialaphos. These experiments demonstrated that the *Sau*3A fragment could be subcloned without loss of activity. They also provided for the proper orientation as shown in the fig. 2, enclosed thereafter. The protein encoded by these clones was detected by using coupled transcription-translation systems derived from extracts of *S. lividans* (ref. 7). Depending on the orientation of the *Sau*3A fragment, translation products of different sizes were observed; 22 Kd for pBG94 and \pm 28 Kd for pBG93. This indicated that the *Sau*3A fragment did not contain the entire resistance gene and that a fusion protein was formed which included a polypeptide sequence resulting from the translation of a pUC19 sequence.

In order to obtain large amounts of the protein, a 1.7 Kb *Pst*I fragment from pBG1 was cloned into the high copy number Streptomycete replicon pIJ385. The obtained plasmid, pBG20, was used to transform *S. hygroscopicus*. Transformants which contained this plasmid had more than 5 times as much PPT acetylating activity and also had increased amounts of a 22 kd protein on sodium dodecylsulfate gels (SDS gels). Furthermore, both the acetyl transferase and the 22 kd protein appeared when the production of Bialaphos began. The correlation of the *in vitro* data, kinetics of synthesis, and amplified expression associated with pBG20 transformants strongly implied that this 22 Kd band was the gene product.

The complete nucleotide sequence of the 625 bp *Sau*3A fragment was determined as well as of flanking sequences. Computer analysis revealed the presence of an open reading frame over the entire length of the *Sau*3A fragment.

Characterization of the sfr gene product

A series of experiments were performed to determine that the open reading frame of the "sfr" gene indeed encoded the Bialaphos resistance enzyme. To determine the 5' end of the resistance gene, the NH₂-terminal sequence of the enzyme was determined. As concerns more particularly the technique used to determine the said sequence, reference is made to the technique developed by J. VANDEKERCKHOVE, Eur. J. Biochem. 152, p. 9-19, 1985, and to French patent applications n° 85 14579 filed on October 1st, 1985 and n° 85 13046 filed on September 2nd, 1985, all of which are incorporated herein by reference.

This technique allows the immobilization on glass fibre sheets coated with the polyquaternary amine commercially available under the registered trademark POLYBRENE of proteins and of nucleic acids previously separated on a sodium dodecylsulfate containing polyacrylamide gel. The transfer is carried out essentially as for the protein blotting on nitrocellulose membranes (ref. 8). This allows the determination of amino-acid composition and partial sequence of the immobilized proteins. The portion of the sheet carrying the immobilized 22 kd protein produced by *S. hygroscopicus* pBG20 was cut out and the disc was mounted in the reaction chamber of a gas-phase sequencer to subject the glass-fibre bound 22 Kd protein to the Edman degradation procedure. The following amino-acid sequence was obtained:

Pro-Glu-Arg-Arg-Pro-Ala-Asp-Ile-Arg-Arg

This sequence matched an amino-acid sequence which was deduced from the open reading frame of the 625 bp *Sau*3A fragment. It corresponded to the stretch from codon 3 to codon 12.

Thus, the NH₂-terminus of the 22 Kd protein was upstream of this sequence. It was determined that translation of the actual protein was likely to be initiated 2 amino-acids earlier at a GTG initiation codon. GTG is often used as initiator codon in *Streptomyces* and translated as methionine. The protein translated from the GTG initiation codon would be 183 amino-acids long and would have a molecular weight of 20 550. This was in good agreement with the observed approximate molecular weight of 22 000.

Furthermore, the termination codon, TGA, was located just downstream of the *Sau*3A site. Cloning of the 625 bp *Sau*3A fragment in a *Bam*HI site digested pUC19 did not result in the reconstruction of the termination codon. This explained the fusion proteins which were observed in the *in vitro* transcription-translation analysis.

Mechanism of PPT-resistance

Having defined a first phenotype and some of the physical characteristics of the resistance gene and its gene product, a series of experiments was then carried out to understand the mechanism by which it confers resistance. As described hereabove, PPT is the portion of Bialaphos which inhibits glutamine synthetase (GS) and that N-acetyl PPT is not an inhibitor. Using a standard assay (ref. 9), *S. hygrosopicus* ATCC 21 705 derivatives were shown to contain a PPT acetyl transferase which was not found in *S. lividans*. The activity does not acetylate the Bialaphos tripeptide. *S. lividans* carrying the resistance gene cloned in pBG20 or pBG16 (a plasmid containing the 625 bp Sau3A fragment cloned into another streptomycete vector, pIJ680) also contained the activity which could acetylate PPT but not Bialaphos. The PPT derived reaction product produced by extracts of pBG20/*S. lividans* was isolated in order to confirm that it was indeed acetyl-PPT. Analysis by mass spectroscopy showed that the molecular weight had increased relative to PPT by the equivalent of one acetyl group. It was thus concluded that the 625 bp Sau3A fragment contained sequences which code for PPT acetyl transferase.

The experimental conditions and reagents used in the techniques disclosed hereabove were as follows :

Preparation and composition of the mediums and buffers above used

1° P medium : 10.3 g of sucrose, 0.025 g of K₂SO₄, 0.203 g of MgCl₂.6H₂O and 0.2 ml of a trace element solution are dissolved in 80 ml of distilled water and autoclaved. Then in order, 1 ml of KH₂PO₄ (0.5%), 10 ml of CaCl₂.2H₂O (3.68 %), and 10 ml of TES buffer (0.25 M), pH : 7.2) are added. Trace element solution (per litre) : ZnCl₂, 40 mg ; FeCl₃.6H₂O, 200 mg ; CuCl₂.2H₂O, 10 mg ; MnCl₂.4H₂O, 10 mg ; Na₂B₄O₇.10H₂O, 10 mg ; (NH₄)₆Mo₇O₂₄.4H₂O, 10 mg.

2° R2YE : 10.3 g of sucrose, 0.025 g of K₂SO₄, 1.012 g of MgCl₂.6H₂O, 1 g of glucose, 0.01 g of Difco casamino acids, and 2.2 g of Difco agar are dissolved in 80 ml distilled water and autoclaved. 0.2 ml of trace element solution, 1 ml of KH₂PO₄ (0.5%), 8.02 ml of CaCl₂.2H₂O (3.68 %), 1.5 ml of L-proline (20 %), 10 ml of TES buffer (0.25 M) (pH : 7.2), 0.5 ml of (1 M) NaOH, 5 ml of yeast extract (10 %) are sequentially added.

3° TE : 10 mM TRIS HCl, 1 mM EDTA, pH 8.0.

4° YEME : Difco yeast extract (0.3 %), Difco peptone (0.5 %), oxoid malt extract (0.3 %), glucose (1 %).

Transformation of *S. lividans* protoplasts

1. A culture composed of 25 ml YEME, 34 % sucrose, 0.005 M MgCl₂, 0.5 % glycine, in a 250 ml baffled flask, is centrifuged during 30 to 36 hours.

2. The pellet is suspended in 10.3 % sucrose and centrifuged. This washing is repeated once.

3. The mycelium is suspended in 4 ml lysozyme solution (1 mg/ml in P medium with CaCl₂ and MgCl₂ concentrations reduced to 0.0025 M) and incubated at 30° C for 15 to 60 minutes.

4. The solution is mixed by pipetting three times in a 5 ml pipette and incubated for further 15 minutes.

5. P medium (5 ml) is added and mixed by pipetting as in step 4.

6. The solution is filtered through cotton wool and protoplasts are gently sedimented in a bench centrifuge at 800 x G during 7 minutes.

7. Protoplasts are suspended in 4 ml P medium and centrifuged again.

8. Step 7 is repeated and protoplasts are suspended in the drop of P medium left after pouring off the supernatant (for transformation).

9. DNA is added in less than 20 µl TE.

10. 0.5 ml PEG 1 000 solution (2.5 g PEG dissolved in 7.5 ml of 2.5% sucrose, 0.0014 K₂SO₄, 0.1 M CaCl₂, 0.05 M TRIS-maleic acid, pH 8.0, plus trace elements) is immediately added and pipetted once to mix the components.

11. After 60 seconds, 5 ml of P medium are added and the protoplasts are sedimented by gentle centrifugation.

12. The pellet is suspended in P medium (1 ml).

13. 0.1 ml is plated out on R2YE plates (for transformation dry plates to 85% of their fresh weight e. g. in a laminar flow cabinet).

14. Incubation at 30° C.

A - Construction of a "sfr" gene cassette

A "sfr" gene cassette was constructed to allow subsequent cloning in plant expression vectors.

Isolation of a FokI-BglII fragment from the plasmid pBG39 containing a "sfr" gene fragment led to the loss of the first codons, including the initiation codon, and of the last codons, including the stop codon.

This fragment of the "sfr" gene could be reconstructed in vitro with synthetic oligonucleotides which encode appropriate amino-acids.

The complementary synthetic oligonucleotides were 5'-CATGAGCCCAGAAC and 3'-TCGGGTCTTGCTGC. By using such synthetic oligonucleotides, the 5' end of the "sfr" gene could be reformed and the GTG initiation codon substituted for a codon well translated by plant cells, particularly an ATG codon.

The DNA fragment containing the oligonucleotides linked to the "sfr" gene was then inserted into an

appropriate plasmid, which contained a determined nucleotide sequences thereafter designated by an "adapter" fragment.

This adapter fragment comprised :

- a TGA termination codon which enabled the last codons of the "sfr" gene to be reformed ;
- appropriate restriction sites which enabled the insertion of the fragment of the nucleotide sequence comprising the "sfr" gene partially reformed with the synthetic oligonucleotides ; this insertion resulted in the reconstruction of an intact "sfr" gene ;
- appropriate restriction sites for the isolation of the entire "sfr" gene.

The "sfr" gene was then inserted into another plasmid, which contained a suitable plant promoter sequence. The plant promoter sequence consisted of the cauliflower mosaic virus promoter sequence (p35S). Of course the invention is not limited to the use of this particular promoter. Other sequences could be chosen as promoters suitable in plants, for example the TR 1'-2' promoter region and the promoter fragment of a Rubisco small subunit gene from *Arabidopsis thaliana* hereafter described.

1° Construction of the plasmid pLK56.2 (fig. 3)

The construction of plasmid pLK56.2 aimed at obtaining a suitable adaptor including the following sequence of restriction sites : SmaI, BamHI, NcoI, KpnI, BglII, MluI, BamHI, HindIII and XbaI.

The starting plasmids used for this construction, pLK56, pJB64 and pLK33 were those disclosed by BOTTERMAN (ref. 11).

The DNA fragments hereafter described were isolated and separated from low melting point agarose (LGA). The plasmid pLK56 was cleaved by the enzymes BamHI and NdeI. A NcoI-NdeI fragment (referred to in the drawings by arc "a" in broken line) obtained from plasmid pJB64 was substituted in pLK56 for the BamHI-NdeI fragment shown at "b". Ligation was possible after filling in the BamHI and NcoI protruding ends with the DNA polymerase I of *E. coli* (Klenow's fragment).

Particularly recircularization took place by means of a T4 DNA ligase. A new plasmid pLK56.3 was obtained. This plasmid was cleaved by the enzymes XbaI and PstI.

The BamHI-PstI fragment of pLK33 (c) (on fig. 3) was substituted for the XbaI-PstI fragment (d) of pLK56.3, after repairing of their respective ends by Klenow's fragment.

After recircularization by means of the T4 DNA ligase, the obtained plasmid pLK56.2 contained a nucleotide sequence which comprised the necessary restriction sites for the subsequent insertion of the "sfr" gene.

2° Construction of the plasmid pGSH150 (fig. 4A)

Parallel with the last discussed construction, there was produced a plasmid containing a promoter sequence recognized by the polymerases of plant cells and including suitable restriction sites, downstream of said promoter sequence in the direction of transcription, which restriction sites are then intended to enable the accomodation of the "sfr" gene then obtainable from pLK56.2, under the control of said plant promoter.

Plasmid pGV825 is described in DEBLAERE et al. (ref. 10). Plasmid pJB63 is from BOTTERMAN (ref. 11). pGV825 was linearized with PvuII and recircularized by the T4 DNA ligase, resulting in the deletion of an internal PvuII fragment shown at (e), (plasmid pGV956).

pGV956 was then cleaved by BamHI and BglII. The BamHI-BglII fragment (f) obtained from pJB63 was dephosphorylated with calf intestine phosphatase (CIP) and substituted for the BamHI-BglII fragment of pGV956.

Plasmid pGV1500 was obtained after recircularization by means of T4 DNA ligase. An EcoRI-HindIII fragment obtained from plasmid pGSH50 was purified. The latter plasmid carried the dual TR 1'-2' promoter fragment described in VELTEN et al., (ref.13). This fragment was inserted in pGV1500, digested with HpaI and HindIII and yielded pGSH150.

This plasmid contains the promoter fragment in front of the 3' end of the T-DNA transcript 7 and a BamHI and ClaI sites for cloning.

3° Construction of the plasmid pGSJ260 (fig. 4B)

CP3 is a plasmid derived from pBR322 and which contains the 35S promoter region of cauliflower mosaic virus within a BamHI fragment.

pGSH150 was cut by BamHI and BglII.

The BamHI-BglII fragment (h) of CP3, which contained the nucleotide sequence of p35S promoter, was substituted for the BamHI-BglII fragment (i) in pGSH150 to form plasmid pGSJ250. pGSJ250 was then opened at its BglII restriction site.

A BamHI fragment obtained from mGV2 (ref. 12) was inserted in pGSJ250 at the BglII site to form plasmid pGSJ260.

However prior to inserting the "sfr" gene obtainable from pLK56.2 into plasmid pGSJ260, it was still desirable to further modify the first in order to permit insertion in a more practical manner. Thus pLK56.2 was further modified as discussed below to yield pGSR1.

Starting from plasmid pGSJ260, two plasmid constructions for subsequent transformations of plant cells were made :

- a first plasmid permitting the expression of the "sfr" gene in the cytoplasm of plant cells, and
- a second plasmid so modified as to achieve transport of the Bialaphos-resistance enzymes to the

chloroplasts of plant cells.

First case : plasmid enabling the expression of the "sfr" gene in the cytoplasm of plant cells

Cloning of the sfr gene cassette in a plant expression vector (pGSR2) (fig. 5)

On figure 5A, the nucleotide sequence of the adapter of pLK56.2 is shown. In particular, the locations of BamHI, NcoI, BglII restriction sites are shown.

This adapter fragment was cleaved by the enzymes NcoI and BglII.

Figure 5B shows the FokI-BglII fragment (j) obtained from pBG39. The locations of these two restriction sites are shown on figure 2.

Using synthetic oligonucleotides, the first codons of the "sfr" gene were reformed, particularly the 5' end of the gene in which a ATG initiation codon was substituted for the initial GTG codon.

This FokI-BglII fragment completed with the synthetic oligonucleotides was then substituted in pLK56.2 for the NcoI-BglII fragment of the adapter. The 3' end of the gene was thus reformed too, after recircularization with T4 DNA ligase. The plasmid obtained, pGSR1, thus contained the entire "sfr" gene inserted in its adapter.

The plasmid pGSJ260 was then opened by BamHI (fig. 5C) and the BamHI fragment obtained from pGSR1, which contained the entire "sfr" gene, was inserted into pGSJ260.

The obtained plasmid, pGSR2 (see fig. 6A) contained a pBR322 replicon, a bacterial streptomycin resistance gene (SDM-SP-AD-transferase) and an engineered T-DNA consisting of :

- the border fragments of the T-DNA ;
- a chimeric kanamycin gene which provided a dominant selectable marker in plant cells ; and
- a chimeric "sfr" gene.

The chimeric "sfr" gene consisting of :

- the promoter region of the cauliflower mosaic virus (p35S) ;
- the "sfr" gene cassette as described in fig. 5 ;
- the 3' untranslated region, including the polyadenylation signal of T-DNA transcript 7.

pGSR2 was introduced into *Agrobacterium tumefaciens* recipient C58CIRif^R (pGV2260) according to the procedure described by DEBLAERE et al. (ref. 10).

This strain was used to introduce the chimeric "sfr" gene in *N. tabacum* SR₁ plants.

Two variant plasmids deriving from pGSR2, namely pGSFR280 and pGSFR281, have been constructed. They differ in the untranslated sequence following the transcription initiation site. In pGSR2, this fragment consists of the following sequence :

GAGGACACGCTGAAATCACCAGTCTCGGATCCATG ;

while it consists of :

GAGGACACGCTGAAATCACCAGTCTCTCTACAAATCGATCCATG

in pGSFR280 and of

GAGGACACGCTGAAATCACCAGTCTCTCTACAAATCGATG

in pGSFR281, with an ATG codon being the initiation codon of the "sfr" gene. The "sfr" gene is also fused to the TR1'-2' promoter in the plasmid pGSH150 (fig. 4A) yielding pGSFR160 and pGSFR161 (fig. 6B). These plasmids contain slight differences in the pTR2 "sfr" gene configuration : the "sfr" gene is correctly fused to the endogenous gene 2' ATG in pGSFR161 (for sequences see ref. 13), whereas 4 extra base pairs (ATCC) are present just ahead of the ATG codon in pGSFR160. Otherwise, plasmids p65FR161 and p65FR160 are completely identical.

All plasmids are introduced in *Agrobacterium* by cointegration in the acceptor plasmids pGV2260 yielding the respective plasmids pGSFR1280, pGSFR1281, pGSFR1160 and pGSFR1161.

Second case : construction of a plasmid containing the "sfr" gene downstream of a DNA sequence encoding a transit peptide and suitable for achieving subsequent translocation of the "sfr" gene expression product into plant-cell-chloroplasts

In another set of experiments, the nucleotide sequence which contained the "sfr" gene was fused to a DNA sequence encoding a transit peptide so as to enable its transport into chloroplasts.

A fragment of the "sfr" gene was isolated from the adapter fragment above described and fused to a transit peptide. With synthetic oligonucleotides, the entire "sfr" gene was reconstructed and fused to a transit peptide.

The plasmid (plasmid pATS3 mentioned below) which contained the nucleotide sequence encoding the transit peptide comprised also the promoter sequence thereof.

Construction of the plasmid pGSR4 which contains the "sfr" gene fused to a DNA sequence encoding transit peptide (fig. 7)

Plasmid pLK57 is from BOTTERMAN, (ref.11). Plasmid pATS3 is a pUC19 clone which contains a 2Kb *EcoRI* genomic DNA fragment from *Arabidopsis thaliana* comprising the promoter region and the transit peptide nucleotide sequence of the gene, the expression thereof is the small subunit of ribulose biphosphate carboxylase (ssu). The *A. thaliana* small subunit was isolated as a 1 500 bp *EcoRI*-*SphI* fragment. The *SphI* cleavage site exactly occurs at the site where the coding region of the mature ssu protein starts.

Plasmids pLK57 and pATS3 were opened with EcoRI and SphI. After recircularization by means of the T4 DNA ligase, a recombinant plasmid pLKAB1 containing the sequence encoding the transit peptide (Tp) and its promoter region (Pssu) was obtained.

In order to correctly fuse the "sfr" gene at the cleavage site of the signal peptide, the N-terminal gene sequence was first modified. Since it was observed that N-terminal gene fusions with the "sfr" gene retain their enzymatic activity, the second codon (AGC) was modified to a GAC, yielding an NcoI site overlapping with the ATG initiator site. A new plasmid, pGSSFR2 was obtained. It only differs from pGSR1 (fig. 5B), by that mutation. The NcoI-BamHI fragment obtained from pGSSFR2 was fused at the SphI end of the transit peptide sequence. In parallel, the "sfr" gene fragment was fused correctly to the ATG initiator of the ssu gene (not shown in figures).

Introduction of the "sfr" gene into a different plant species

The Bialaphos-resistance induced in plants by the expression of chimeric genes, when the latter have been transformed with appropriate vectors containing said chimeric genes, has been demonstrated as follows. The recombinant plasmids containing the "sfr" gene were introduced separately by mobilization into Agrobacterium strain C58C₁ Rif^R (pGV2260) according to the procedure described by DEBLAERE and al., Nucl. Acid. Res., 13, p. 1 477, 1985. Recombinant strains containing hybrid Ti plasmides were formed. These strains were used to infect and transform leaf discs of different plant species, according to a method essentially as described by HORSH and al., 1985, Science, vol. 227. Transformation procedure of these different plant species given by way of example, is described thereafter.

1. Leaf disc transformation of Nicotiana tabacum

Used Media are described thereafter :

A₁ MS salt/2 + 1% sucrose
0.8% agar
pH 5.7

A₁₀ B5-medium + 250 mg/l NH₄NO₃
750 mg/l CaCl₂ 2H₂O
0.5 g/l 2-(N-Morpholino)ethane-
sulfonic acid (MES) pH 5.7
30 g/l sucrose

A₁₁ B5-medium + 250 mg/l NH₄NO₃
0.5 g/l MES pH 5.7
2 % glucose
0.8 % agar
40 mg/l adenine
+ 1 mg/l 6-Benzylaminopurine
(BAP)

0.1 mg/l Indole-3-acetic acid
(IAA)

500 mg/l Claforan

A₁₂ B5-medium + 250 mg/l NH₄NO₃
0.5 g/l MES pH 5.7
2 % glucose
0.8 % agar

40 mg/l adenine
+ 1 mg/l BAP

200 mg/l claforan

A₁₃ MS-salt/2 + 3 % sucrose

0.5 MES g/l pH 5.7

0.7 % agar

200 mg/l claforan

Bacterial medium = min A : (Miller 1972) 60 mM

K₂HPO₄, 3H₂O,

33 mM KH₂PO₄ ; 75 mM (NH₄)₂SO₄

1.7 M trisodiumcitrat; 1 mM MgSO₄ ;

2 g/l glucose ; 50 mg/l vitamine B1

- Plant material :

Nicotiana tabacum cv. Petit Havana SR1

Plants are used 6 to 8 weeks after subculture on medium A₁

- Infection :

- midribs and edges are removed from leaves.

- Remaining parts are cut into segments of about 0.25 cm² and are placed in the infection medium A₁₀ (about

12 segments in a 9 cm Petri dish containing 10 ml A₁₀).

- Segments are then infected with 25 µl per Petri dish of a late log culture of the Agrobacterium strain grown in min A medium.

- Petri dish are incubated for 2 to 3 days at low light intensity.

5 - After 2 to 3 days medium is removed and replaced by 20 ml of medium A₁₀ containing 500 mg/l clarofan.

- Selection and shoot induction

- The leaf discs are placed on medium A₁₁ containing a selective agent :

100 mg/l kanamycin and

10 to 100 mg/l phosphinotricin.

- Leaf discs are transferred to fresh medium weekly.

- After 3 to 4 weeks regenerating calli arise. They are separated and placed on medium A₁₂ with the same concentration of selective agent as used for the selection.

15 - Rooting

- After 2 to 3 weeks the calli are covered with shoots, which can be isolated and transferred to rooting medium A₁₃ (without selection).

- Rooting takes 1 to 2 weeks.

- After a few more weeks, these plants are propagated on medium A₁.

20

2. Tuber disc infection of Solanum tuberosum (potato)

Used media are described thereafter :

C₁ B5-medium + 250 mg/l NH₄NO₃

300 mg/l (CaCH₂PO₄)₂

25 0.5 g/l MES pH 5.7

0.5 g/l polyvinylpyrrolidone (PVP)

40 g/l mannitol (= 0.22M)

0.8% agar

40 mg/l adenine

30 C₂ B5-medium + 250 mg/l NH₄NO₃

400 mg/l glutamine

0.5 g/l MES pH 5.7

0.5 g/l PVP

40 g/l mannitol

35 40 mg/l adenine

0.8 % agar

+ 0.5 mg/l transzeatine

0.1 mg/l IAA

500 mg/l clarofan

40 C₅ MS salt/2 + 3 % sucrose

0.7 % agar

pH 5.7

C₇ B5-medium + 250 mg/l NH₄NO₃

400 mg/l glutamine

45 0.5 g/l MES pH 5.7

0.5 g/l PVP

20 g/l mannitol

20 g/l glucose

40 mg/l adenine

50 0.6 % agarose

+ 0.5 mg/l transzeatine

0.1 mg/l IAA

500 mg/l clarofan

C₈ B5-medium + 250 mg/l NH₄NO₃

55 400 mg/l glutamine

0.5 g/l MES pH 5.7

0.5 g/l PVP

20 g/l mannitol

20 g/l glucose

50 40 mg/l adenine

0.6 % agarose

+ 200 mg/l clarofan

1 mg/l transzeatine

C₉ B5-medium + 250 mg/l NH₄NO₃

55 400 mg/l glutamine

0.5 g/l MES pH 5.7	
0.5 g/l PVP	
20 g/l mannitol	
20 g/l glucose	
40 mg/l adenine	
0.6 % agarose	
+ 1 mg/l transzeatine	5
0.01 mg/l Gibberellic acid A ₃ (GA ₃)	
100 mg/l clarofan	
C ₁₁ MS salt/2 + 6 % sucrose	10
0.7 % agar	
Bacterial medium = min A : (Miller 1972 60 mM K ₂ HPO ₄ ·3H ₂ O;	
33 mM KH ₂ PO ₄ ; 7.5 mM (NH ₄) ₂ SO ₄ ;	
1.7 trisodiumcitrat; 1 mM MgSO ₄ ;	
2 g/l glucose; 50 mg/l vitamines B ₁ .	15
- <u>Plant material</u>	
Tubers of <u>Solanum tuberosum</u> c.v. Berolina	
c.v. Désirée	20
- <u>Infection</u>	
- Potatoes are peeled and washed with water.	
- Then they are washed with concentrated commercial bleach for 20 minutes, and	
- rinsed 3 to 5 times with sterile water.	
- The outer layer is removed (1 to 1.5 cm)	25
- The central part is cut into discs of about 1 cm ² and 2 to 3 mm thick.	
- Discs are placed on medium C ₁ (4 pieces in a 9 cm Petri dish).	
- 10 µl of a late log culture of an <u>Agrobacterium</u> strain grown in min A medium is applied on each disc.	
- Discs are incubated for 2 days at low light intensity.	30
- <u>Selection and shoot induction</u>	
- Discs are dried on a filter paper and transferred to medium C ₂ with 100 mg/l kanamycin.	
- After one month small calli are removed from the discs and transferred to medium C ₇ containing 50 mg/l kanamycin.	35
- After a few more weeks, the calli are transferred to medium C ₈ containing 50 mg/l kanamycin.	
- If little shoots start to develop, the calli are transferred to elongation medium C ₉ containing 50 mg/l kanamycin.	
- <u>Rooting</u>	
- Elongated shoots are separated and transferred to rooting medium C ₁₁ .	40
- Rooted shoots are propagated on medium C ₅ .	
3. <u>Leaf disc infection of Lycopersicum esculentum</u> (tomato)	
Used media are described thereafter	
A ₁ MS salt/2 + 1 % sucrose	45
0.8 % agar	
pH 5.7	
B ₁ B5-medium + 250 mg/l NH ₄ NO ₃	
0.5 g/l MES pH 5.7	
0.5 g/l PVP	50
300 mg/l Ca (H ₂ PO ₄) ₂	
2 % glucose	
40 mg/l adenine	
40 g/l mannitol	
B ₂ B5-medium + 250 mg/l NH ₄ NO ₃	55
0.5 g/l MES pH 5.7	
0.5 g/l PVP	
400 mg/l glutamine	
2 % glucose	
0.6 % agarose	60
40 mg/l adenine	
40 g/l mannitol	
+ 0.5 mg/l transzeatine	
0.01 mg/l IAA	65

- 500 mg/l claforan
 B₃ B5-medium + 250 mg/l NH₄NO₃
 0.5 g/l MES pH 5.7
 0.5 g/l PVP
 5 400 mg/l glutamine
 2 % glucose
 0.6 % agarose
 40 mg/l adenine
 30 g/l mannitol
 10 + 0.5 mg/l transzeatine
 0.01 mg/l IAA
 500 mg/l clarofan
 B₄ B5-medium + 250 mg/l NH₄NO₃
 0.5 g/l MES pH 5.7
 15 0.5 g/l PVP
 400 mg/l glutamine
 2 % glucose
 0.6 % agarose
 40 mg/l adenine
 20 20 g/l mannitol
 + 0.5 mg/l transzeatine
 0.01 mg/l IAA
 500 mg/l clarofan
 B₅ B5-medium + 250 mg/l NH₄NO₃
 25 0.5 g/l MES pH 5.7
 0.5 g/l PVP
 400 mg/l glutamine
 2 % glucose
 0.6 % agarose
 30 40 mg/l adenine
 10 g/l mannitol
 + 0.5 mg/l transzeatine
 0.01 mg/l IAA
 500 mg/l clarofan
 35 B₆ B5-medium + 250 mg/l NH₄NO₃
 0.5 g/l MES pH 5.7
 0.5 g/l PVP
 400 mg/l glutamine
 2 % glucose
 40 0.6 % agarose
 40 mg/l adenine
 + 0.5 mg/l transzeatine
 0.01 mg/l IAA
 200 mg/l clarofan
 45 B₇ B5-medium + 250 mg/l NH₄NO₃
 0.5 g/l MES pH 5.7
 0.5 g/l PVP
 400 mg/l glutamine
 2 % glucose
 50 0.6 % agarose
 40 mg/l adenine
 + 1 mg/l transzeatine
 200 mg/l clarofan
 B₈ MS salt/2 + 2 % sucrose
 55 0.5 g/l MES pH 5.7
 0.7 % agar
 B₉ B5-medium + 250 mg/l NH₄NO₃
 0.5 g/l MES pH 5.7
 0.5 g/l PVP
 60 2 % glucose
 0.6 % agarose
 40 mg/l adenine
 + 1 mg/l transzeatine
 0.01 mg/l GA₃
 65 Bacterial medium = min A : (Miller 1972) 60 mM

K₂HPO₄·3H₂O ;
 33 mM KH₂PO₄; 7.5 mM (NH₄)₂SO₄;
 1.7 M trisodiumcitrat; 1 mM MgSO₄ ;
 2 g/l glucose; 50 mg/l vitamine B1

- Plant material

Lycopersicum esculentum cv. Lucullus.

Plants are used 6 weeks after subculture on medium A₁.

- Infection

- Midrib is removed from the leaves.
- Leaves are cut in segments of about 0.25 to 1 cm² (the edges of the leaves are not wounded, so that only maximum 3 sides of the leaf pieces is wounded).
- Segments are placed in infection medium B₁ (upside down), about 10 segments in a 9 cm Petri dish.
- Segments are then infected with 20 µl per Petri dish of a late log culture of the *Agrobacterium* strain grown in min A medium.
- Petri dishes incubate for 2 days at low light intensity.
- Medium is removed after 2 days and replaced by 20 ml of medium B₁ containing 500 mg/l clarofan.

- Selection and shoot induction

- The leaf discs are placed in medium B₂ + 50 or 100 mg/l kanamycin.
- Each 5 days the osmotic pressure of the medium is lowered by decreasing the mannitol concentration, transfers are done consecutively in medium B₃, B₄, B₅, and B₆.
- After one month calli with meristems are separated from the leaf discs and placed on medium B₇ with 50 or 100 mg/l kanamycin.
- Once little shoots have formed, calli are transferred to elongation medium B₉ with 50 or 100 mg/l kanamycin.

- Rooting

- Elongated shoots are separated and transferred to medium B₈ for rooting.
- Plants are propagated on medium A₁.

Greenhouse tests for herbicide resistance

Material and method

In this experiment, two herbicides comprising phosphinotricin as active ingredient, are used.

These compounds are those commercially available under the registered trademarks BASTAR[®] and MEIJI HERBIACER.

These products are diluted to 2 % with tap water. Spraying is carried out on a square metre area from the four corners. Temperature of the greenhouse is about 22°C for tobaccos and tomato, and above 10°C to 15°C for potato.

Results

- Tobacco spraytest

a) *Nicotiana tabacum* cv. Petit Havana SR1 plants transformed with the chimeric "sfr" genes as present in pGSFR1161 or pGSFR1281, as well as untransformed control plants (from 10 cm to 50 cm high) are treated with 20 l BASTAR/ha. Control SR1 plants die after 6 days, while transformed plants are fully resistant to 20 l BASTAR/ha and continue growing undistinguishable from untreated plants. No visible damage is detected, also the treatment is repeated every two weeks. The treatment has no effect in subsequent flowering. The recommended dose of BASTAR herbicide in agriculture is 2.5-7.5 l/ha.

b) A similar experiment is performed using 8 l/ha MEIJI HERBIACER. The transformed plants (the same as above) are fully resistant and continue growing undistinguishable from untreated plants. No visible damage is detectable.

- Potato spraytest

Untransformed and transformed potato plants (*Solanum tuberosum* cv. Berolina) (20 cm high) with the chimeric "sfr" gene as present in pGSFR1161 or pGSFR1281 are treated with 20 l BASTAR/ha. Control plants die after 6 days while the transformed plants do not show any visible damage. They grow undistinguishable from untreated plants.

- tomato spraytest

Untransformed and transformed tomato plants (*lycopersium esculentum* c.v. luculus) (25 cm high) with the chimeric "sfr" gene as present in pGSFR1161 and pGSFR1281 are treated with 20 l BASTAR/ha. control plants die after six days while transformed plants are fully resistant. They do not show any visible damage and grow undistinguishable from untreated plants.

- Growth control of phytopathogenic fungi with transformed plants

In another set of experiments, potato plants expressing chimeric "sfr" genes as present in pGSFR1161 or pGSFR1281 are grown in a greenhouse compartment at 20°C under high humidity. Plants are inoculated by spraying 1 ml of a suspension of 10^6 *Phytophthora infestans* spores per ml. Plants grow in growth chambers (20°C, 95 % humidity, 4 000 lux) until fungal disease symptoms are visible (one week). One set of the plants are at that moment sprayed with Bialaphos at a dose of 8 l/ha. Two weeks later, untreated plants are completely ingested by the fungus. The growth of the fungus is stopped on the Bialaphos treated plants and no further disease symptoms evolve. The plants are effectively protected by the fungicide action of Bialaphos.

- Transmission of the PPT resistance through seeds

Transformed tobacco plants expressing the chimeric "sfr" gene present in pGSFR1161 and pGSFR1281 are brought to flowering in the greenhouse. They show a normal fertility.

About 500 F1 seeds of each plant are sown in soil, F1 designating seeds of the first generation, i.e directly issued from the originally transformed plants. When seedlings are 2-3 cm high, they are sprayed with 8 l BASTAR/ha. 7 days later, healthy and damaged plants can be distinguished in a ratio of approximately 3 to 1. This shows that PPT resistance is inherited as a dominant marker encoded by a single locus.

10 resistant F1 seedlings are grown to maturity and seeds are harvested. F2 seedlings are grown as described above and tested for PPT-resistance by spraying BASTAR at a dose of 8 l/ha. Some of the F1 plants produce F2 seedlings which are all PPT-resistant showing that these plants are homozygous for the resistance gene. The invention also concerns plant cells and plants non-essentially-biologically-transformed with a GS inhibitor-inactivating-gene according to the invention.

In a preferred embodiment of the invention, plant cells and plants are non-biologically-transformed with the "sfr" gene hereabove described.

Such plant cells and plants possess, stably integrated in their genome, a non-variety-specific character which render them able to produce detectable amounts of phosphinotricin-acetyl transferase.

This character confers to the transformed plant cells and plants a non-variety-specific enzymatic activity capable of inactivating or neutralizing GS inhibitors like Bialaphos and PPT.

Accordingly, plant cells and plants transformed according to the invention are rendered resistant against the herbicidal effects of Bialaphos and related compounds.

Since Bialaphos was first described as a fungicide, transformed plants can also be protected against fungal diseases by spraying with the compound several times.

In a preferred embodiment, Bialaphos or related compounds is applied several times, particularly at time intervals of about 20 to 100 days.

The invention also concerns a new process for selectively protecting a plant species against fungal diseases and selectively destroying weeds in a field comprising the steps of treating a field with an herbicide, wherein the plant species contain in their genome a DNA fragment encoding a protein having an enzymatic activity capable of neutralizing or inactivating GS inhibitors and wherein the used herbicide comprises as active ingredient a GS inhibitor.

It comes without saying that the process according to the invention can be employed with the same efficiency, either to only destroy weeds in a field, if plants are not infected with fungi, either to only stop the development of fungi if the latter appears after destruction of weeds.

In a preferred embodiment of the process according to the invention, plant species are transformed with a DNA fragment comprising the "sfr" gene as described hereabove, and the used herbicide is PPT or a related compound.

Accordingly, a solution of PPT or related compound is applied over the field, for example by spraying, several times after emergence of the plant species to be cultivated, until early and late germinating weeds are destroyed.

It is quite evident that before emergence of plant species to be cultivated, the field can be treated with an herbicidal composition to destroy weeds.

On the same hand, fields can be treated even before the plant species to be cultivated are sowed. Before emergence of the desired plant species, fields can be treated with any available herbicide, including Bialaphos-type herbicides.

After emergence of the desired plant species, Bialaphos or related compound is applied several times. In a preferred embodiment, the herbicide is applied at time intervals of about from 20 to 100 days.

Since plants to be cultivated are transformed in such a way as to resist to the herbicidal effects of Bialaphos-type herbicides, fields can be treated even after emergence of the cultivated plants.

This is particularly useful to totally destroy early and late germinating weeds, without any effect on the plants to be produced.

Preferably, Bialaphos or related compound is applied at a dose ranging from about 0.4 to about 1.6 kg/ha, and diluted in a liquid carrier at a concentration such as to enable its application to the field at a rate ranging from about 2 to about 8 l/ha.

There follows examples, given by way of illustration, of some embodiments of the process with different plant species.

- Sugarbeets

The North European sugarbeet is planted from March 15 up to April 15, depending upon the weather condition and more precisely on the precipitation and average temperature. the weeds problems are more or less the same in each country and can cause difficulties until the crop closes its canopy around mid-July.

Weed problems can be separated in three situations :

- early germination of the grassy weeds,
- early germinating broadleaved weeds,
- late germinating broadleaved weeds.

Up to now, pre-emergence herbicides have been successfully used. Such compounds are for example those commercially available under the registered trademarks : PYRAMIN^R, GOLTIX^R and VENZARR^R. However, the susceptibility to dry weather conditions of these products as well as the lack of residual activity to control late germinating weeds have led the farmer to use post-emergence products in addition to pre-emergence ones.

Table (I) thereafter indicates the active ingredients contained in the herbicidal compositions cited in the following examples.

TABLE (I)

Commercial Name	Active Ingredient	Formulation	
AVADEX ^R	Diallate	EC 400 g/l	
AVADEX BW ^R	Triallate	EC 400 g/l	25
GOLTIX ^R	Metamitron	WP 70 %	
RONEET ^R	Cycloate	EC 718 g/l	
TRAMAT ^R	Ethofumerate	EC 200 g/l	30
FERVINAL ^R	Alloxydime-sodium	SP 75 %	
BASTA ^R	Phosphinotricin	200 g/l	
PYRAMIN FL ^R	Chloridazon	SC 430 g/l	35

According to the invention, post-emergence herbicides consist of Bialaphos or related compounds, which offer a good level of growth control of annual grasses (Bromus, Avena spp., Alopecurus, POA) and broadleaves (Galium, Polygonum, Senecio, Solanum, Mercurialis).

Post-emergence herbicides can be applied at different moments of the growth of sugarbeet ; at a cotyledon level, two-leave level or at a four-leave level.

Table (II) thereafter represents possible systems of field-treatment, given by way of example.

In those examples, the post-emergence herbicide of the class of Bialaphos used is BASTAR^R, in combination with different pre-emergence herbicides. Concentrations are indicated in l/ha or kg/ha.

TABLE (II)

POSSIBLE WEEDCONTROL SYSTEMS IN SUGARBEETS, BASED ON THE USE OF BASTA^R, PROVIDING BEETS ARE MADE RESISTANT AGAINST THE LATTER CHEMICAL (in lt or kg/ha).

Pre-sowing	Pre-emergence	Cotyledons	Two-leaves	Four leaves
1. AVADEx ^R 3.5 lt	-	BASTA ^R 3 lt	BASTA ^R /Lramat 3 lt 1.5 lt	-
2. AVADEx ^R 3.5 lt	GOLTIX ^R 4 kg	-	-	-
3. RONEET ^R 4 lt	GOLTIX ^R 5 kg	-	-	-
4. RONEET ^R 4 lt	GOLTIX ^R 2.5 kg	-	BASTA ^R 3 lt	-
5. TRAMAT ^R 5 lt	-	-	BASTA ^R 3 lt	BASTA ^R /GOLTIX ^R 2 lt 2 kg
6. -	GOLTIX ^R 2.5 kg	-	BASTA ^R 3 lt	-
7. -	-	BASTA ^R /Lramat 3 lt 1.7 lt	-	BASTA ^R /GOLTIX ^R 3 lt 2 kg
8. PYRAMIN ^R 6 lt	-	BASTA ^R 3 lt	Venzar 1 kg	-
9. -	-	BASTA ^R 3 lt	BASTA ^R /GOLTIX ^R 3 lt 2 kg	-
10. DIALATE ^R 3.5 lt	PYRAMIN ^R 6 lt	-	BASTA ^R /Metamitron 3 lt 1 kg	-

- Potatoes

Potatoes are grown in Europe on about 8.10⁶ Ha. The major products used for weed control are Linuron/monolinuron or the compound commercially available under the denomination METRABUZIN. These products perform well against most weedspecies.

However, weeds such as Galium and Solanum plus late germinating Chenopodium and Polygonum are not always effectively controlled, while control of the annual grasses is also sometime erratic.

Once again, late germinating broadleaved weeds are only controllable by post-emergence applications of herbicides such as BASTAR.

Table (III) thereafter represents some examples given by way of example of field-treatment in the case of potatoes.

TABLE (III)

Weeds control systems in potatoes, based on the use of BASTAR^R, providing potatoes are rendered resistant to BASTAR^R.

Linuron + monolinuron (375 g + 375 g/ha) prior to emergence

BASTAR^R 3-4 lt/ha after emergence (5-15 cm)

BASTAR^R/fluazifop-butyl 3-4 lt/ha + 2 lt/ha after emergence (5-15 cm)

Linuron

WP 50 % (AFALON^R)

Monolinuron

WP 47.5 % (ARESSIN^R)

fluazifop-butyl

EL 250 g/l (FUSILADE^R)

The strains pGSJ260 and pBG39 used hereabove have been deposited on December 12nd, 1985, at the "German Collection of Micro-organisms" (DEUTSCHE SAMMLUNG VON MIKROORGANISMEN) at Göttingen, Germany. They received the deposition numbers DSM 3 606 and DSM 3 607 respectively.

Further embodiments of the invention are described hereafter with reference to the figures in which :

- fig. 8 shows the restriction map of a plasmid pJS1 containing another Bialaphos-resistance-gene ;

- fig. 9 shows the nucleotide sequence of the "sfrsv" gene containing the resistance gene ;

- fig. 10 shows the amino acid homology of "sfrsv" gene and "sfr" gene,

- fig. 11 shows the construction of a plasmid, given by way of example, which contains the "sfrsv" gene and suitable for the transformation of plant cells.

Another Bialaphos-resistance-gene has been isolated from another Bialaphos-producing-strains, i.e. streptomyces viridochromogenes. This second resistance-gene is thereafter designated by "sfrsv" gene.

This second preferred DNA fragment according to the invention, for the subsequent transformation of plant cells, consists of a nucleotide sequence coding for at least part of a polypeptide having the following sequence

V S P E R R P V E I R P A T A A D M
 5 A A V C D I V N H Y I E T S T V N F
 10 R T E P Q T P Q E W I D D L E R L Q
 15 D R Y P W L V A E V E G V V A G I A
 20 Y A G P W K A R N A Y D W T V E S T
 25 V Y V S H R H Q R L G L G S T L Y T
 30 H L L K S M E A Q G F K S V V A V I
 35 G L P N D P S V R L H E A L G Y T A
 40 F W Q R D F E L P A P P R P V R P V
 45 T Q I *

50 which part of said polypeptide is of sufficient length to confer protection against Bialaphos-"plant-protecting-capability"-, to plant cells, when incorporated genetically and expressed therein. Reference will also be made here-after to the "plant-protecting-capability" against Bialaphos of the abovesaid nucleotide sequence.

Meaning of the designation of amino acids by a single letter is given thereafter.

Alanine A
 Arginine R
 Asparagine N
 55 Aspartic Acid D
 Cysteine C
 Cystine C
 Glycine G
 Glutamic Acid E
 60 Glutamine Q
 Histidine H
 Isoleucine I
 Leucine L
 Lysine K
 65 Methionine M

p
 S.
 (fi
 hy
 a r
 fra
 we.

Phenylalanine F
Proline P
Serine S
Threonine T
Tryptophan W
Tyrosine Y
Valine V

This second preferred DNA fragment consists of the following nucleotide sequence :

TAAAGAGGTGCCCCGCCACCCGCTTTCGCAGAACACCGAAGGAGACCACAC
↓
GTGAGCCCAGAACGACGCCCGGTGAGATCCGTCCCGCCACCGCCGCCGA
CATGGCGGCGGTCTGCGACATCGTCAATCACTACATCGAGACGAGCACGG
TCAACTTCCGTACGGAGCCGCAGACTCCGCAGGAGTGGATCGACGACCTG
GAGCGCCTCCAGGACCGCTACCCCTGGCTCGTCGCCGAGGTGGAGGGCGT
CGTCGCCGGCATCGCCTACGCCGGCCCCCTGGAAGGCCCGCAACGCCTACG
ACTGGACCGTCGAGTCGACGGTGTACGTCTCCACCGGCACCAGCGGCTC
GGACTGGGCTCCACCCTCTACACCCACCTGCTGAAGTCCATGGAGGCCCA
GGGCTTCAAGAGCGTGGTTCGCCGTCATCGGACTGCCCAACGACCCGAGCG
TGCGCCTGCACGAGGCGCTCGGATACACCGCGCGCGGGACGCTGCGGGCA
GCCGGCTACAAGCACGGGGGCTGGCACGACGTGGGGTTCTGGCAGCGCGA
CTTCGAGCTGCCGGCCCCGCCCCGCCCCGTCCGGCCCGTCACACAGATCT
↓
GAGCGGAGAGCGCATGGC

5

10

15

20

25

30

35

40

45

50

55

60

65

or of a part thereof expressing a polypeptide having plant-protecting capability against Bialaphos ;

There follows hereafter the description of experiments carried out for the isolation of the "sfrsv" resistance gene, the construction of expression vectors which contain the resistance gene and which allow the subsequent transformation of plant cells, in order to render them resistant GS inhibitors.

Cloning of the bialaphos-resistance-"sfrsv" gene from *Streptomyces viridochromogenes*

The strain *Streptomyces viridochromogenes* DSM 40736 (ref 1) was grown and total DNA of this strain was prepared according to standard techniques. DNA samples were digested respectively with *Pst*I, *Sma*I and *Sau*3A1 in three different reactions and separated on an agarose gel, together with plasmid DNA from pGSR1 (fig. 5B) digested with *Bam*HI. In a Southern analysis the DNA was blotted on a nitrocellulose filter and hybridized with the labeled *Bam*HI fragment from pGSR1 containing the "sfr" gene. In all four lanes of the gel, a restriction fragment was showing strong homology with the probe : a *Pst*I fragment of about 3 kb, a *Sma*I fragment of about 1.2 kb and *Sau*3A1 fragment of 0.5 kb. In order to clone this gene, *Pst*I restriction fragments were directly cloned in the *Escherichia coli* vector pUC8. 3000 colonies obtained after transformation were

transferred to nitrocellulose filters, and hybridized with the "sfr" probe. Positive candidates were further tested for their growth on minimal medium plates containing 300 µg/ml PPT. One transformant that grew on PPT-containing-medium was further analysed. The plasmid map and relevant restriction sites of this plasmid pJS1 are represented in fig. 8. The strain MC1061 (pJS1) has been deposited on March 06, 1987 at the DEUTSCHE SAMMLUNG VON MIKROORGANISMEN (DSM) under deposition number DSM 4023. The clone restriction fragment has been sequenced according to the Maxam and Gilbert method and the coding region of the gene could be identified through homology. The sequence of the "sfrsv" gene is represented in fig.9 and the homology on the nucleotide and amino acid sequence level with "sfr" gene is shown in fig. 10.

Expression of the "sfrsv" gene

A "sfrsv gene cassette" was also constructed to allow subsequent cloning in plant expression vectors. A *Ban*II-*Bgl*III fragment containing the "sfrsv" coding region without the initiation codon GTG was isolated from pJS1. This fragment was ligated in the vector pLK56-2 digested with *Nco*I and *Bgl*III, together with a synthetic oligonucleotide 5'-CATGAGCC-3', similar with the one described for "sfr" gene and shown in fig. 5. The construction of pGSR1SV is schematically shown in fig. 11. Since similar cassettes of both genes are present in respectively pGSR1 and pGSR1SV, previous described constructions for the expression of the "sfr" gene in plants can be repeated.

Enzymatic analysis of crude extracts from *E. coli* strains carrying plasmid pGSR1SV demonstrated the synthesis of an acetylase which could acetylate PPT. This was shown by thin layer chromatography of the reaction products.

The "sfrsv" gene was then inserted into the plasmid vector pGSJ260 (fig. 4B) under the control of the CaMV 35s promoter, to yield a plasmid pGS2SV, similar to pGSR2 (fig. 6A) except that the "sfrsv" gene is substituted for the "sfr" gene.

It is clear that herbicide resistance genes of the above type may be obtained from many other microorganisms that produce PPT or PPT derivatives. Herbicide resistance gene can then be incorporated in plant cells with a view of protecting them against the action of such Glutamine Synthetase inhibitors. For instance, a Bialaphos-resistance-gene is obtained from *Kitasatosporia* (ref. 15).

Transformed plant cells and plants which contain the "sfrsv" resistance gene can be obtained with plasmid pGSR2SV, using the same *Agrobacterium*-mediated-transformation system as hereabove described for the transformation of different plant species with the "sfr" gene.

Plants are regenerated and tested for their resistance with similar spraying tests as described hereabove. All plants behaved similarly and show resistance against herbicides consisting of glutamine synthetase inhibitors.

Finally, the invention also pertains to the combination of the plants resistant to an inhibitor of glutamine synthetase as defined above with the corresponding inhibitor of glutamine synthetase for use in the production of cultures of said plants free from weeds.

REFERENCES

1. BAYER et al., HELVETICA CHIMICA ACTA, 1972
2. WAKABAYASHI K. and MATSUNAKA S., Proc. 1982, British Crop Protection Conference, 439-450
3. M. MASON et al., PHYTOCHEMISTRY, 1982, vol. 21, no 4, p. 855-857.
4. C. J. THOMPSON et al., NATURE, July 31, 1980, vol. 286, no 5 772, p. 525-527
5. C. J. THOMPSON et al., JOURNAL OF BACTERIOLOGY, August 1982, p. 678-685
6. C. J. THOMPSON et al., GENE 20, 1982, p. 51-62
7. C. J. THOMPSON et al., MOL. GEN. GENET., 1984, 195, p. 39-43
8. TOWBIN et al., PROC. NATL. ACAD. SCI. USA, 1979, 76, p. 4 350-4 354
9. METHODS OF ENZYMOLOGY, V.XLIII, p. 737-755
10. DEBLAERE H. et al., 1985, Nucl. Acid. Res., 13, 1 477
11. BOTTERMAN J., February 1986, Ph. D. Thesis, State University of Ghent
12. DEBLAERE R., february 1986, Ph. D Thesis, Free University of Brussel, Belgium
13. VELTEN et al, EMBO J. 1984, vol. 3, no12, p. 2 723-2 730
14. CHATER et al, Gene cloning in Streptomyces. Curr. Top. Microbiol. Immunol., 1982, 96, p. 69-75
15. OMURA et al, J. of Antibiotics, Vol. 37, 8, 939-940, 1984
16. MURAKAMI et al, Mol. Gen. Genet., 205, 42-50, 1986
17. MANDERSCHIED and WILD, J. Plant Phys., 123, 135-142, 1986

Claims

1. Process for controlling the action in plant cells and plants of a glutamine synthetase inhibitor when contacted herewith, which comprises providing said plants with a heterologous DNA including a foreign nucleotide sequence, capable of being expressed in the form of a protein in said plant cells and plants, under conditions such as to cause said heterologous DNA to be integrated stably through generations in the cells of said plants, and wherein said protein has an enzymatic activity capable of causing inactivation

or neutralization of said glutamine synthetase inhibitor.

2. Process according to claim 1, wherein the heterologous DNA fragment comprises a foreign nucleotide sequence coding for a polypeptide having a PPT acetyl transferase activity.

3. Process according to the claims 1 to 2, wherein the heterologous DNA fragment belongs to the genome of a *Streptomyces*.

4. Process according to any of the claims 1 to 3, wherein the heterologous DNA fragment comprises a foreign nucleotide sequence coding for the following amino acid sequence :

5

10

15

20

25

30

35

40

45

50

55

60

65

183
ARG ARG PRO ALA ASP ILE ARG ARG ALA THR GLU ALA ASP MET PRO

228
ADA VAL CYS THR ILE VAL ASN HIS TYR ILE GLU THR SER THR VAL

273
ASN PHE ARG THR GLU PRO GLN GLU PRO GLN GLU TRP THR ASP ASP

318
LEU VAL ARG LEU ARG GLU ARG TYR PRO TRP LEU VAL ALA GLU VAL

363
ASP GLY GLU VAL ALA GLY ILE ALA TYR ALA GLY PRO TRP LYS ALA

408
ARG ASN ALA TYR ASP TRP THR ALA GLU SER THR VAL TYR VAL SER

453
PRO ARG HIS GLN ARG THR GLY LEU GLY SER THR LEU TYR THR HIS

498
LEU LEU LYS SER LEU GLU ALA GLN GLY PHE LYS SER VAL VAL ALA

543
VAL ILE GLY LEU PRO ASN ASP PRO SER VAL ARG MET HIS GLU ALA

588
LEU GLY TYR ALA PRO ARG GLY MET LEU ARG ALA ALA GLY PHE LYS

633
HIS GLY ASN TRP HIS ASP VAL GLY PHE TRP GLN LEU ASP PHE SER

678
LEU PRO VAL PRO PRO ARG PRO VAL LEU PRO VAL THR GLU ILE

723

in which X represents MET or VAL, or a part of a polypeptide of sufficient length to possess said enzymatic activity and, when said DNA fragment is expressed in plant cells, to protect the latter against the herbicidal activity of glutamine synthetase inhibitors.

5. Process according to any of the claims 1 to 4, wherein the heterologous DNA fragment comprises the following nucleotide sequence :

GTG AGC CCA GAA

123

5 CSA CGC CCG GCC GAC ATC CSC CGT GCC ACC GAG GCG GAC ATG CCG
223

10 GCG GTC TGC ACC ATC GTC AAC CAC TAC ATC GAG ACA AGC ACG GTC
273

AAC TTC CGT ACC GAG CCG CAG GAA CCG CAG GAG TGG ACG GAC GAC
318

15 CTC GTC CGT CTG CCG GAG CCG TAT CCC TGG CTC GTC GCC GAG GTG
363

20 GAC GGC GAG GTC GCC GGC ATC GCC TAC GCG GGC CCC TGG AAG GCA
408

CGC AAC GCC TAC GAC TGG ACG GCC GAG TCG ACC GTG TAC GTC TCC
453

25 CCC CGC CAC CAG CCG ACG GGA CTG GGC TCC ACG CTC TAC ACC CAC
498

30 CTG CTG AAG TCC CTG GAG GCA CAG GGC TTC AAG AGC GTG GTC GGT
543

GTC ATC GGG CTG CCC AAC GAC CCG AGC GTG CCG ATG CAC GAG GCG
588

35 CTC GGA TAT GCC CCC CCG GGC ATG CTG CCG GCG GCC GGC TTC AAG
633

40 CAC GGG AAC TGG CAT GAC GTG GGT TTC TGG CAG CTG GAC TTC AGC
678

45 CTG CCG GTA CCG CCC CGT CCG GTC CTG CCC GTC ACC GAG ATC
723

or a part thereof expressing a polypeptide capable of inactivating said glutamine synthetase inhibitor.

6. Process according to any of claims 1 to 5, wherein the initiation codon ATG is substituted for the initiation codon GTG of the sequence defined in claim 5.

7. Process according to any of claims 1 to 3, wherein the heterologous DNA fragment comprises a foreign nucleotide sequence coding for the following amino acid sequence :

0 242 246

V S P E R R P V E I R P A T A A D M	
A A V C D I V N H Y I E T S T V N F	5
R T E P Q T P Q E W I D D L E R L Q	10
D R Y P W L V A E V E G V V A G I A	15
Y A G P W K A R N A Y D W T V E S T	20
V Y V S H R H Q R L G L G S T L Y T	25
H L L K S M E A Q G F K S V V A V I	30
G L P N D P S V R L H E A L G Y T A	35
R G T L R A A G Y K H G G W H D V G	40
F W Q R D F E L P A P P R P V R P V	45
T Q I *	

or a part of a polypeptide of sufficient length to possess said enzymatic activity and, when said DNA fragment is expressed in plant cells, to protect the latter against the herbicidal activity of glutamine synthetase inhibitors.

8. Process according to claim 7, wherein the heterologous DNA fragment comprises the following nucleotide sequence :

TAAAGAGGTGCCCCGCCACCCGCTTTCGCAGAACACCGAAGGAGACCACAC
 ↓
 5 GTGAGCCCAGAACGACGCCCGGTTCGAGATCCGTCCCGCCACCGCCGCCGA
 CATGGCGGCGGTCTGCGACATCGTCAATCACTACATCGAGACGAGCACGG
 10 TCAACTTCCGTACGGAGCCGCAGACTCCGCAGGAGTGGATCGACGACCTG
 15 GAGCGCCTCCAGGACCGCTACCCCTGGCTCGTCGCCGAGGTGGAGGGCGT
 CGTCGCCGGCATCGCCTACGCCGGCCCCCTGGAAGGCCCGCAACGCCTACG
 20 ACTGGACCGTCGAGTCGACGGTGTACGTCTCCACCGGCACCAGCGGCTC
 25 GGACTGGGCTCCACCCTCTACACCCACCTGCTGAAGTCCATGGAGGCCCA
 30 1 GGGCTTCAAGAGCGTGGTCCCGTCATCGGACTGCCCAACGACCCGAGCG
 TCGCCTGCACGAGGCGCTCGGATACACCGCGCGCGGGACGCTGCGGGCA
 35 GCCGGCTACAAGCACGGGGGCTGGCACGACGTGGGGTTCTGGCAGCGCGA
 CTTTCGAGCTGCCGGCCCCCGCCCCGCCCCGTCCGGCCCGTCACACAGATCT
 40 GAGCGGAGAGCGCATGGC ↑

or a part thereof expressing a polypeptide capable of inactivating said glutamine synthetase inhibitor.

45 9. In a process for producing plants and reproduction material of said plants including a heterologous genetic material stably integrated therein and capable of being expressed in said plants or reproduction material in the form of a protein capable of inactivating or neutralizing the activity of a glutamine synthetase inhibitor, comprising the non biological steps of producing plants cells or plant tissue including said heterologous genetic material from starting plant cells or plant tissue not able to express that inhibiting or neutralizing activity, regenerating plants or reproduction material of said plants or both from said plant cells or plant tissue including said genetic material and, optionally, biologically replicating said last mentioned plants or reproduction material or both, wherein said steps of producing said plant cells or plant tissue including said heterologous genetic material comprises transforming said starting plant cells or plant tissue with a DNA recombinant containing a nucleotide sequence encoding said protein as well as the regulatory elements selected among those which are capable of enabling the expression of said nucleotide sequence in said plant cells or plant tissue and to cause the stable integration of said nucleotide sequence in said plant cells and tissue, as well as in the plant and reproduction material processed therefrom throughout generations.

60 10. The process according to claim 9, wherein starting cells are transformed with a recombinant DNA which contains the fragment of any one of the claims 1 to 8 recombined with heterologous DNA, and wherein said recombinant DNA is devoid of any DNA region involved in the biosynthesis of Bialaphos, when said fragment originates itself from a Bialaphos-producing strain.

65 11. The process according to claim 10, wherein the recombinant DNA is a vector suitable for the transformation of a cellular host and, when introduced therein, for allowing for the neutralization or inhibition of glutamine synthetase inhibitors.

12. The process according to claim 11, wherein the vector comprises said recombinant DNA under the control of replicon elements suitable for the transformation of bacteria, particularly *E. coli*.
13. The process of claim 11, wherein the vector comprises said recombinant DNA fragment under the control of a plant promoter region, which vector further comprises regulation elements allowing for the expression of said DNA fragments in plant cells, when the latter are later transformed with said vector.
14. The process of any of claims 10 to 13, which confers resistance against herbicidal effects of Bialaphos, PPT or related derivatives to the transformed plant cells. 5
15. The process of claim 13 or 14, wherein the vector comprises a nucleotide sequence encoding a transit peptide intercalated between said plant promoter region and said recombinant DNA fragment.
16. The process of claim 15, wherein the transit peptide is selected from ribulose-1,5 biphosphate carboxylase and chlorophyll a/b binding proteins. 10
17. The process of any of claims 11 to 16, wherein said vector is a Ti plasmid.
18. Plant cells, non biologically transformed, wherein they possess, stably integrated in their genome, a DNA fragment encoding a protein having a non-variety-specific enzymatic activity capable of neutralizing or inactivating glutamine synthetase inhibitors.
19. Plant cells according to claim 18, wherein they are capable of generating a plant capable of producing seeds, said seeds having a non-variety-specific enzymatic activity capable of inactivating or neutralizing glutamine synthetase inhibitors. 15
20. Plant cells according to claim 18 or 19, wherein they are transformed by the process of any of claims 9 to 17.
21. Plant cells according to any of claims 18 to 20, which produce detectable amounts of phosphinotricin acetyl transferase. 20
22. Seeds, non biologically transformed, which possess, stably integrated in their genome, a DNA fragment encoding a protein having a non-variety-specific enzymatic activity capable of inactivating or neutralizing glutamine synthetase inhibitors.
23. Seeds according to claim 22, which are capable of germinating into a plant capable of producing seeds having a non-variety-specific enzymatic activity capable of inactivating or neutralizing glutamine synthetase inhibitors. 25
24. Seeds according to claim 22 or 23, which are transformed by the process of any one of the claims 9 to 17.
25. Plants, non biologically transformed, which possess, stably integrated in their genome, a DNA fragment encoding a protein having a non-variety-specific enzymatic activity capable of neutralizing or inactivating glutamine synthetase inhibitors. 30
26. Plants according to claim 25, which are capable of producing seeds having a non-variety specific enzymatic activity capable of inactivating or neutralizing glutamine synthetase inhibitors.
27. Plants according to claim 25 or 26, which are transformed by the process of any of claims 9 to 17. 35
28. Process for protecting a plant species and selectively destroying weeds in a field comprising the steps of treating a field with an herbicide, wherein the plant species contain in their genome a DNA fragment encoding a protein having an enzymatic activity capable of neutralizing or inactivating glutamine synthetase inhibitors, and wherein the used herbicide is a glutamine synthetase inhibitor.
29. Process according to claim 28, wherein the plant species contains a DNA fragment according to any one of claims 1 to 8. 40
30. Process according to claim 28 or 29, wherein the plant species are transformed according to the process of any one of claims 9 to 17.
31. Process according to any one of claims 28 to 30, wherein a solution of a glutamine synthetase inhibitor is applied on the field, after emergence of the cultivated plant species, several times, particularly at time intervals of about 20 to 100 days, until early and late germinating weeds are destroyed. 45
32. Process according to any of claims 28 to 31, wherein glutamine synthetase inhibitors comprise Bialaphos, phosphinotricin and related compounds.
33. Process for selectively protecting a plant species in a field against fungal diseases comprising the steps of treating a field with an herbicide, wherein the plant species contain in their genome a DNA fragment encoding a protein having an enzymatic activity capable of neutralizing an inactivating glutamine synthetase inhibitors and wherein the used herbicide is a glutamine synthetase inhibitor. 50
34. Process according to claim 33, wherein the plant species contain a DNA fragment according to any of claims 1 to 8.
35. Process according to claim 33 or 34, wherein the plant species are transformed according to the process of any one of claims 9 to 17. 55
36. Process according to any one of claims 33 to 35, wherein a solution of a glutamine synthetase inhibitor is applied on the field, after emergence of the cultivated plant species, several times, particularly at time intervals of about 20 to 100 days until fungi are destroyed.
37. Process according to any one of claims 33 to 36, wherein glutamine synthetase inhibitor is selected from a group which comprises Bialaphos, phosphinotricin and related compounds. 60
38. Process according to claim 32 or 37, wherein Bialaphos, PPT or related compound is applied at a dose ranging from about 0.4 to about 1.6 kg/ha.
39. Process according to claim 38, wherein Bialaphos, PPT or related compounds is diluted in a liquid 65

carrier at a concentration such as to enable its application to the field at a rate ranging from about 2 l/ha to about 8 l/ha.

40. Process according to any of claim 38 or 39, wherein selectively protected plant species comprise sugar-beet, rice, potato, tomato, maize, tobacco.

41. Vector which contains a DNA fragment encoding a polypeptide having an enzymatic activity capable of causing inactivation of glutamine synthetase inhibitors.

42. Vector according to claim 41, which is suitable for the transformation of plant cells and plants.

43. Vector according to claim 41 or 42, which contains a DNA fragment belonging to the genome of a Streptomyces.

44. Vector according to any of claims 41 to 43, which contains the DNA fragment recombined with heterologous DNA, wherein said recombinant DNA is devoid of any DNA region involved in the biosynthesis of Bialaphos.

45. The vector of claim 44 suitable for the transformation of a cellular host and, when introduced therein, for allowing for expression of a polypeptide capable of inactivating glutamine synthetase inhibitor.

46. The vector of claim 45, wherein said DNA fragment is under the control of replicon elements suitable for the transformation of bacteria, particularly E. coli.

47. The vector of claim 46, wherein said DNA fragment is under the control of a plant promoter region, and of regulation elements allowing for the expression of said DNA fragments in plant cells, when the latter are later transformed with said vector.

48. The vector of claim 47, which confers bialaphos-resistance to the transformed plant cells.

49. The vector of claim 48, wherein a sequence encoding a transit peptide is intercalated between said plant promoter region and said DNA fragment.

50. The vector of claim 49, wherein the transit peptide is selected from ribulose-1,5 biphosphate carboxylase and chlorophyl a/b binding proteins.

51. The vector of any of claims 45 to 50, wherein said vector is a modified Ti plasmid.

52. Vector according to any of claims 41 to 51, which comprises the following nucleotide sequence :

193 . GTG AGC CCA GAA

CGA CGC CCG GCC GAC ATC CGC CGT GCC ACC GAG GCG GAC ATG CCG 5
229

GCG GTC TGC ACC ATC GTC AAC CAC TAC ATC GAG ACA AGC ACB GTC
273

AAC TTC CGT ACC GAG CCG CAG GAA CCG CAG GAG TGG ACG GAC GAC 10
318

CTC GTC CGT CTG CCG GAG CGC TAT CCC TGG CTC GTC GCC GAG GTG 15
363

GAC GGC GAG GTC GCC GGC ATC GCC TAC GCG GGC CCC TGG AAG GCA
408

CGC AAC GCC TAC GAC TGG ACG GCC GAG TCG ACC GTG TAC GTC TCC 20
453

CCC CGC CAC CAG CCG ACG GGA CTG GGC TCC ACG CTC TAC ACC CAC 25
499

CTG CTG AAG TCC CTG GAG GCA CAG GGC TTC AAG AGC GTG GTC GCT
543

GTC ATC GGG CTG CCC AAC GAC CCG AGC GTG CGC ATG CAC GAG GCG 30
589

CTC GGA TAT GCC CCC CGC GGC ATG CTG CCG GCG GCC GGC TTC AAG 35
633

CAC GGG AAC TGG CAT GAC GTG GGT TTC TGG CAG CTG GAC TTC AGC
678

CTG CCG GTA CCG CCC CGT CCB GTC CTG CCC GTC ACC GAG ATC 40
723

53. The vector of claim 52, wherein the initiation codon ATG is substituted for the initiation codon GTG of the sequence defined in claim 52. 45

54. Vector according to any of claims 41 to 51, which comprises the following nucleotide sequence : 50

50

55

60

65

TAAAGAGGTGCCCCGCCACCCGCTTTCGCAGAACACCGAAGGAGACCACAC

↓

5 GTGAGCCCAGAACGACGCCCGGTCGAGATCCGTCCCGCCACCGCCGCCGA

CATGGCGGCGGTCTGCGACATCGTCAATCACTACATCGAGACGAGCACGG

10 TCAACTTCCGTACGGAGCCGCAGACTCCGCAGGAGTGGATCGACGACCTG

15 GAGCGCCTCCAGGACCGCTACCCCTGGCTCGTCGCCGAGGTGGAGGGCGT

CGTCGCCGGCATCGCCTACGCCGGCCCCCTGGAAGGCCCCGCAACGCCTACG

20 ACTGGACCGTCGAGTCGACGGTGTACGTCTCCACCGGCACCAGCGGCTC

25 GGACTGGGCTCCACCCTCTACACCCACCTGCTGAAGTCCATGGAGGCCCA

GGGCTTCAAGAGCGTGGTCGCCGTCATCGGACTGCCCAACGACCCGAGCG

30 TCGCCTGCACGAGGCGCTCGGATACACCGCGCGCGGGACGCTGCGGGCA

35 GCCGGCTACAAGCACGGGGGCTGGCACGACGTGGGGTTCTGGCAGCGCGA

CTTCGAGCTGCCGGCCCCCGCCCCGCCCCGTCCGGCCCGTCACACAGATCT

↑

40 GAGCGGAGAGCGCATGGC

45 55. A DNA fragment, for the subsequent transformation of plant cells, coding for a polypeptide having phosphinothricin-acetyl-transferase activity, which consists of a nucleotide sequence coding for the following amino acid sequence :

50

55

60

65

X SER PRO GLU

123
 ARG ARG PRO ALA ASP ILE ARG ARG ALA THR GLU ALA ASP MET PRO 5

229
 ALA VAL CYS THR ILE VAL ASN HIS TYR ILE GLU THR SER THR VAL

273
 ASN PHE ARG THR GLU PRO GLN GLU PRO GLN GLU TRP THR ASP ASP 10

318
 LEU VAL ARG LEU ARG GLU ARG TYR PRO TRP LEU VAL ALA GLU VAL 15

363
 ASP GLY GLU VAL ALA GLY ILE ALA TYR ALA GLY PRO TRP LYS ALA

408
 ARG ASN ALA TYR ASP TRP THR ALA GLU SER THR VAL TYR VAL SER 20

453
 PRO ARG HIS GLN ARG THR GLY LEU GLY SER THR LEU TYR THR HIS 25

498
 LEU LEU LYS SER LEU GLU ALA GLN GLY PHE LYS SER VAL VAL ALA

543
 VAL ILE GLY LEU PRO ASN ASP PRO SER VAL ARG MET HIS GLU ALA 30

588
 LEU GLY TYR ALA PRO ARG GLY MET LEU ARG ALA ALA GLY PHE LYS 35

633
 HIS GLY ASN TRP HIS ASP VAL GLY PHE TRP GLN LEU ASP PHE SER

678
 LEU PRO VAL PRO PRO ARG PRO VAL LEU PRO VAL THR GLU ILE 40

723 45

in which X represents MET or VAL, or of a part of said nucleotide sequence of sufficient length to code for a polypeptide still having phosphinothricin-acetyl-transferase activity

56. The DNA fragment of claim 55 which comprises the following nucleotide sequence :

50

55

60

65

123 GTG AGC CCA GAA

5 CGA CGC CCG GCC GAC ATC CGC CGT GCC ACC GAG GCG GAC ATG CCG
228

GCG GTC TGC ACC ATC GTC AAC CAC TAC ATC GAG ACA AGC ACG GTC
273

10 AAC TTC CGT ACC GAG CCG CAG GAA CCG CAG GAG TGG ACG GAC GAC
318

15 CTC GTC CGT CTG CCG GAG CCG TAT CCC TGG CTC GTC GCC GAG GTG
363

GAC GGC GAG GTC GCC GGC ATC GCC TAC GCG GGC CCC TGG AAG GCA
408

20 CGC AAC GCC TAC GAC TGG ACG GCC GAG TCG ACC GTG TAC GTC TCC
453

25 CCC CGC CAC CAG CCG ACG GGA CTG GGC TCC ACG CTC TAC ACC CAC
498

CTG CTG AAG TCC CTG GAG GCA CAG GGC TTC AAG AGC GTG GTC GCT
543

30 GTC ATC GGG CTG CCC AAC GAC CCG AGC GTG CCG ATG CAC GAG GCG
583

35 CTC GGA TAT GCC CCC CCG GGC ATG CTG CCG GCG GCC GGC TTC AAG
633

CAC GGG AAC TGG CAT GAC GTG GGT TTC TGG CAG CTG GAC TTC AGC
678

40 CTG CCG GTA CCG CCC CGT CCG GTC CTG CCC GTC ACC GAG ATC
723

45

or a part thereof expressing a polypeptide having phosphinotricin-acetyl-transferase activity.

57. A DNA fragment, for the subsequent transformation of plant cells, coding for a polypeptide having phosphinotricin-acetyl-transferase activity, which consists of a nucleotide sequence coding for the following amino acid sequence :

50

55

60

65

0 242 246

V S P E R R P V E I R P A T A A D M	
A A V C D I V N H Y I E T S T V N F	5
R T E P Q T P Q E W I D D L E R L Q	10
D R Y P W L V A E V E G V V A G I A	15
Y A G P W K A R N A Y D W T V E S T	20
V Y V S H R H Q R L G L G S T L Y T	25
H L L K S M E A Q G F K S V V A V I	30
G L P N D P S V R L H E A L G Y T A	35
R G T L R A A G Y K H G G W H D V G	40
F W Q R D F E L P A P P R P V R P V	45
T Q I *	50

or a part of said nucleotide sequence of sufficient length to code for a polypeptide still having phosphinotricin-acetyl-transferase activity.

58. The DNA fragment of claim 57, which comprises the following nucleotide sequence :

55

60

65

TAAAGAGGTGCCCCGCCACCCGCTTTCGCAGAACACCGAAGGAGACCACAC

↓

5 GTGAGCCCAGAACGACGCCCGGTCGAGATCCGTCCCGCCACCGCCGCCGA

CATGGCGGCGGTCTGCGACATCGTCAATCACTACATCGAGACGAGCACGG

10

TCAACTTCCGTACGGAGCCGCAGACTCCGCAGGAGTGGATCGACGACCTG

15

GAGCGCCTCCAGGACCGCTACCCCTGGCTCGTCGCCGAGGTGGAGGGCGT

CGTCGCCGGCATCGCCTACGCCGGCCCCCTGGAAGGCCCGCAACGCCTACG

20

ACTGGACCGTCGAGTCGACGGTGTACGTCTCCACCGGCACCAGCGGCTC

25

GGACTGGGCTCCACCCTCTACACCCACCTGCTGAAGTCCATGGAGGCCCA

GGGCTTCAAGAGCGTGGTCGCCGTCATCGGACTGCCCAACGACCCGAGCG

30

TGCGCCTGCACGAGGCGCTCGGATACACCGCGCGCGGGACGCTGCGGGCA

35

GCCGGCTACAAGCACGGGGGCTGGCACGACGTGGGGTTCTGGCAGCGCGA

CTTCGAGCTGCCGGCCCCGCCCCGCCCCGTCCGGCCCGTCACACAGATCT

40

GAGCGGAGAGCGCATGGC

or of a part thereof expressing a polypeptide having phosphinotricin-acetyl-transferase activity.

45

50

55

60

65

0242246




FIG.1

0242246

CCC GCT CAA GCT CGC TGT CAT TTT CGA GAC GCC ATC TTT GGA AGC
GGT GGC CGA ATC CGT ACT GCG CGG ACT CGA CGA CGC GTA AAA CGA
TCG ACC ACG TAC ACG AGT CCG GAC ACG GGG CGA GGA GGC CCG GTT
CCG GCA CCG AGG AAG ACC GAA GGA AGA CCA CAC GTG AGC CCA GAA
CGA CGC CCG GCC GAC ATC CGC CGT GCC ACC GAG GCG GAC ATG CCG
FokI↑
GGG GTC TGC ACC ATC GTC AAC CAC TAC ATC GAG ACA AGC ACG GTC
AAC TTC CGT ACC GAG CCG CAG GAA CCG CAG GAG TGG ACG GAC GAC
CTC GTC CGT CTG CCG GAG CGC TAT CCC TGG CTC GTC GCC GAG GTG
GAC GGC GAG GTC GCC GGC ATC GCC TAC GCG GGC CCC TGG AAG GCA
CGC AAC GCC TAC GAC TGG ACG GCC GAG TCG ACC GTG TAC GTC TCC
CCC CGC CAC CAG CCG ACG GGA CTG GGC TCC ACG CTC TAC ACC CAC
CTG CTG AAG TCC CTG GAG GCA CAG GGC TTC AAG AGC GTG GTC GCT
GTC ATC GGG CTG CCC AAC GAC CCG AGC GTG CGC ATG CAC GAG GCG
CTC GGA TAT GCC CCC CGC GGC ATG CTG CCG GCG GCC GGC TTC AAG
CAC GGG AAC TGG CAT GAC GTG GGT TTC TGG CAG CTG GAC TTC AGC
CTG CCG GTA CCG CCC CGT CCG GTC CTG CCC GTC ACC GAG ATC TGA
ACG GAG TGC GCG TGG GCA TCG CCC GAG TTG GAG CTG GTA CCG GAA
CTC ATC GAA CTC AAC TGG CAT ACC CGC AAT GGT GAG GTG GAA CCG
CGG CCG ATC GCG TAC GAC CGT GCC CAG G

FIG.2

0242246




FIG.3




FIG. 4A

0242246




FIG.4B

0242246





FIG.5A





FIG.5B





FIG.5C





FIG. 6A



- ① LB
- ② PTR1
- ③ PTR2
- ④ 3' END 7
- ⑤ T-DNA
- ⑥ R.B.

FIG. 6B

0242246




FIGURE 8

0242246

TAAAGAGGTGCCCGCCACCCGCTTTCGCAGAACACCGAAGGAGACCACAC
↓
GTGAGCCCAGAACGACGCCCGGTGAGATCCGTCCCGCCACCGCCGCCGA
CATGGCGGGCGGTCTGCGACATCGTCAATCACTACATCGAGACGAGCACGG
TCAACTTCCGTACGGAGCCGCAGACTCCGCAGGAGTGGATCGACGACCTG
GAGCGCCTCCAGGACCGCTACCCCTGGCTCGTCGCCGAGGTGGAGGGCGT
CGTCGCCGGCATCGCCTACGCCGGCCCCCTGGAAGGCCCGCAACGCCTACG
ACTGGACCGTCGAGTCGACGGTGTACGTCTCCCACCGGCACCAGCGGCTC
GGACTGGGCTCCACCCTCTACACCCACCTGCTGAAGTCCATGGAGGCCCA
GGGCTTCAAGAGCGTGGTCGCCGTCATCGGACTGCCCAACGACCCGAGCG
TGCGCCTGCACGAGGCGCTCGGATACACCGCGCGCGGGACGCTGCGGGCA
GCCGGCTACAAGCACGGGGGCTGGCACGACGTGGGGTTCTGGCAGCGCGA
CTTCGAGCTGCCGGCCCCGCCCCGCCCCGTCCGGCCCGTCACACAGATCT
↑
GAGCGGAGAGCGCATGGC

FIGURE 9

0242246

	10	20	30	40	50
PBG39-----	MSPERRPADI	RRATEADMPA	VCTIVNHYIE	TSTVNFRTPE	QEPQEWTDLL
PJS1-----	VSPERRPVEI	RPATAADMAA	VCDIVNHYIE	TSTVNFRTPE	QTPQEWIDDL

	60	70	80	90	100
PBG39-----	VLRLRERYPW	VAEVDGEVAG	IAYAGPWKAR	NAYDWTAESE	VYVSPRHQRT
PJS1-----	ERLQDRYPW	VAEVEGVVAG	IAYAGPWKAR	NAYDWTVEST	VYVSHRHQRL

	110	120	130	140	150
PBG39-----	GLGSTLYTHL	LKSLEAQQFK	SVVAVIGLPN	DPSVRMHEAL	GYAPRGMLRA
PJS1-----	GLGSTLYTHL	LKSMEAQQFK	SVVAVIGLPN	DPSVRLHEAL	GYTARGTLRA

	160	170	180	
PBG39-----	AGFKHGNWHD	VGFWQLDFSL	PVPPRPVLPV	TEI*
PJS1-----	AGYKHGGWHD	VGFWQRDFEL	PAPPRPVRPV	TQI*

FIGURE 10

0242246

pLK562

pJS1




FIGURE 11



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

EP 87 40 0544

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
X	EP-A-0 173 327 (MEIJA SEIKA KAISHA) * Page 19, lines 28-36; figure 7 *	12, 41, 43, 45, 46, 52, 55, 56	C 12 N 15/00 A 01 H 1/00 C 12 N 5/00 A 01 H 5/10 C 12 N 9/10
Y		1-40, 42, 44, 47-51, 53, 54, 57, 58	
Y	WO-A-8 402 920 (MONSANTO) * Page 12, lines 13-18; page 39, lines 1-16; claims 29-34 *	1-58	
Y	JOURNAL OF MOLECULAR AND APPLIED GENETICS, vol. 2, no. 6, 1984, pages 621-635, Raven Press, New York, US; G. DONN et al.: Herbicide-resistant alfalfa cells: An example of gene amplification in plants" * Page 625, right-hand column, last paragraph *	1-58	TECHNICAL FIELDS SEARCHED (Int. Cl. 4) C 12 N A 01 H A 01 N
Y	NATURE, vol. 317, 24th October 1985, page 668; R. SHIELDS: "Engineering herbicide resistance" * Whole article *	1-17, 47-51	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04-07-1987	Examiner MADDOX A.D.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Y	WO-A-8 402 913 (MONSANTO) * Page 4, lines 17-28; page 5, lines 9-20; claims 17-19; page 41; figures 18,21,27,37 *	6,12, 15-17, 47-51, 53	
Y	--- GENE, vol. 33, no. 2, 1985, pages 197-206, Elsevier Science Publishers, Amsterdam, NL; J. VARA et al.: "Cloning and expression of a puromycin N-acetyl transferase gene from Streptomyces alboniger in Streptomyces lividans and Escherichia coli" * Summary; page 204, right-hand column, paragraph 1 *	12,46	
Y	--- BIOTECHNOLOGY, vol. 4, no. 9, September 1986, pages 786-789, New York, US; J.T. FAYERMAN: "New developments in gene cloning in antibiotic producing microorganisms" * Page 788, bottom of left-hand column - top of right-hand column *	7,8,54 ,57,58	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04-07-1987	Examiner MADDOX A.D.

CATEGORY OF CITED DOCUMENTS

- X : particularly relevant if taken alone
Y : particularly relevant if combined with another document of the same category
A : technological background
O : non-written disclosure
P : intermediate document

- T : theory or principle underlying the invention
E : earlier patent document, but published on, or after the filing date
D : document cited in the application
L : document cited for other reasons
& : member of the same patent family, corresponding document



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

EP 87 40 0544

DOCUMENTS CONSIDERED TO BE RELEVANT			Page 3
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Y	HELVETICA CHIMICA ACTA, vol. 55, fasc. 1, no. 25, 1972, pages 224-239; E. BAYER et al.: "Stoffwechselprodukte von Mikroorganismen - Phosphinothricin und Phosphinothricyl-alanyl-alanin" * Summary *	7,8,54 ,57,58	
Y	WO-A-8 602 097 (GENERAL HOSPITAL CORP.) * Page 13, last paragraph - page 14, paragraph 1; page 26; claim 23 *	28-40	
Y	"The Pesticide Manual", 7th edition, page 302, editor C.R. WORTHING, The British Crop Protection Council * Page 302, entry no. 6930 *	28-40	
P,X	CHEMICAL ABSTRACTS, vol. 106, 1987, page 1149, abstract no. 1151u, Columbus, Ohio, US; T. MURAKAMI et al.: "The bialaphos biosynthetic genes of Streptomyces hygroscopicus: molecular cloning and characterization of the gene cluster", & MGG, MOL. GEN. GENET. 1986, 205(1), 42-50 * Abstract *	12,41, 43,45, 46,52, 55,56	
The present search report has been drawn up for all claims			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
Place of search THE HAGUE		Date of completion of the search 04-07-1987	Examiner MADDOX A.D.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

EP 87 40 0544

DOCUMENTS CONSIDERED TO BE RELEVANT

Page 4

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
P, Y	Idem	1-40, 42, 44, 47-51, 53, 54, 57, 58	
A	--- EMBO JOURNAL, vol. 4, 1985, pages 25-32, IRL Press Ltd., Oxford, GB; P.H. SCHREIER et al.: "The use of nuclear-encoded sequences to direct the light-regulated synthesis and transport of a foreign protein into plant chloroplasts" * Page 30, right-hand column - page 31 *	13-17, 47-51	
A	--- PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 81, May 1984, pages 2960-2964; A.R. CASHMORE: "Structure and expression of a pea nuclear gene encoding a chlorophyll a/b-binding polypeptide" * Page 2964 *	13-17, 47-51	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04-07-1987	Examiner MADDOX A.D.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	



DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	CHEMICAL ABSTRACTS, vol. 98, 1983, page 242, abstract no. 48585v, Columbus, Ohio, US; P. LANGELEDDEKE et al.: "Glufosinate (HOE 39866), a new non-selective contact herbicide: results of several years' experimentation in orchards and vineyards from different European countries", & MEDED. FAC. LANDBOUWWET., RIJKSUNIV., GENT 1982, 47(1), 95-104 * Abstract *	31,32,36	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
A	--- CHEMICAL ABSTRACTS, vol. 104, no. 5, February 1986, page 152, abstract no. 29747a, Columbus, Ohio, US; J.D.G. JONES et al.: "High level expression of introduced chimeric genes in regenerated transformed plants", & EMBO J. 1985, 4(10), 2411-18 * Abstract *	16,50	
A	--- NUCLEIC ACIDS RESEARCH, vol. 14, no. 4, February 1986, pages 1565-1581, IRL Press Ltd., Oxford, GB; M. ZALACAIN et al.: "Nucleotide sequence of the hygromycin B phosphotransferase gene from Streptomyces hygrosopicus" * Page 1573, paragraph 1; page 1577, paragraph 2 *	6,53	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04-07-1987	Examiner MADDOX A.D.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			



EP 87 40 0544

Page 6

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
A	NUCLEIC ACIDS RESEARCH, vol. 13, no. 19, October 1985, pages 6981-6998, IRL Press Ltd, Oxford, GB; J. VELTEN et al.: "Selection-expression plasmid vectors for use in genetic transformation of higher plants"		

A	CHEMICAL ABSTRACTS, vol. 104, no. 9, 3rd March 1986, page 311, abstract no. 64619g, Columbus, Ohio, US; M.C. ERICKSON: "Purification and properties of glutamine synthetase from spinach leaves", & PLANT PHYSIOL. 1985, 79(4), 923-7		

P, A	THE JOURNAL OF ANTIBIOTICS, vol. 49, no. 5, May 1986, pages 688-693, Tokyo, JP; T. KOBAYASHI et al.: "Cloning and characterization of the Streptothricin-resistance gene which encodes Streptothricin acetyltransferase from Streptomyces lavendulae"		TECHNICAL FIELDS SEARCHED (Int. Cl.4)

A	GB-A-2 007 976 (MEIJI SEIKA)		

The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 04-07-1987	Examiner MADDOX A.D.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document		T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document	

This Page Blank (uspto)